

MAFK Rabbit pAb

Catalog # AP97487

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	O60675
Other Accession	O60675
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17523
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MAFK
Epitope Specificity	61-156/156
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the bZIP family. Maf subfamily. Contains 1 bZIP domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The developmentally regulated expression of the globin genes depends on upstream regulatory elements termed locus control regions (LCRs). LCRs are associated with powerful enhancer activity that is mediated by the transcription factor NFE2 (nuclear factor erythroid-2). NFE2 recognition sites are also present in the gene promoters of 2 heme biosynthetic enzymes, porphobilinogen deaminase (PBGD; MIM 609806) and ferrochelatase (FECH; MIM 612386). NFE2 DNA-binding activity consists of a heterodimer containing an 18-kD Maf protein (MafF, MafG (MIM 602020), or MafK) and p45 (MIM 601490). Both subunits are members of the activator protein-1 superfamily of basic leucine zipper (bZIP) proteins (see MIM 165160). Maf homodimers suppress transcription at NFE2 sites.[supplied by OMIM, Nov 2008]

Additional Information

Gene ID	7975
Other Names	NFE2U; P18; NF-E2; MAFK_CHICK; MAFK; MAFK_HUMAN; Erythroid transcription factor NF-E2 p18 subunit; MAFK_MOUSE; MAF bZIP transcription factor K; v-maf avian musculoaponeurotic fibrosarcoma oncogene homolog K; basic-leucine zipper transcription factor MafK; nuclear factor erythroid-2, ubiquitous (p18); Transcription factor MafK; NF-E2 ubiquitous subunit
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name MAFK

Function Since they lack a putative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize among themselves (PubMed:[9150357](#)). However, they act as transcriptional activators by dimerizing with other (usually larger) basic-zipper proteins, such as NFE2, NFE2L1/NRF1, NFE2L2/NRF2 and NFE2L3/NRF3, and recruiting them to specific DNA-binding sites (PubMed:[8932385](#), PubMed:[9150357](#)). Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NF-E2 transcription factor (PubMed:[9150357](#)).

Cellular Location Nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.