

FRG1 Rabbit pAb

Catalog # AP97492

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q14331
Other Accession	Q14331
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29172
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FRG1
Epitope Specificity	181-258/258
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the FRG1 family.
SUBUNIT	Identified in the spliceosome C complex.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene maps to a location 100 kb centromeric of the repeat units on chromosome 4q35 which are deleted in facioscapulohumeral muscular dystrophy (FSHD). It is evolutionarily conserved and has related sequences on multiple human chromosomes but DNA sequence analysis did not reveal any homology to known genes. In vivo studies demonstrate the encoded protein is localized to the nucleolus. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	2483
Other Names	FRG1A; FSG1; FRG1_HUMAN; FRG1; FSHD region gene 1 protein; FRG1_MOUSE; FSHD region gene 1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	FRG1 (HGNC:3954)
Function	Binds to mRNA in a sequence-independent manner. May play a role in

regulation of pre-mRNA splicing or in the assembly of rRNA into ribosomal subunits. May be involved in mRNA transport. May be involved in epigenetic regulation of muscle differentiation through regulation of activity of the histone-lysine N-methyltransferase KMT5B.

Cellular Location

Nucleus, Cajal body. Nucleus, nucleolus. Cytoplasm. Cytoplasm, myofibril, sarcomere, Z line. Note=Localization changes during myogenesis from mainly cytoplasmic in undifferentiated myoblasts, to strongly nucleolar in early myotubes and back to cytoplasmic 5 days post-differentiation (PubMed:20970242). Localized at the Z-line in the sarcomere of matured myotubes 8 days post-differentiation (PubMed:20970242).

Tissue Location

Expressed in adult muscle, lymphocytes, fetal brain, muscle, and placenta. Also expressed in the smooth muscle of arteries and veins, the sweat glands and the epidermis

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.