

P2X2 Rabbit pAb

Catalog # AP97562

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9UBL9
Other Accession	Q9UBL9
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51754
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human P2X2
Epitope Specificity	51-150/471
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the P2X receptor family.
SUBUNIT	Functional P2XRs are organized as homomeric and heteromeric trimers.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The product of this gene belongs to the family of purinoceptors for ATP. This receptor functions as a ligand-gated ion channel. Binding to ATP mediates synaptic transmission between neurons and from neurons to smooth muscle. Multiple transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Aug 2013]

Additional Information

Gene ID	22953
Other Names	DFNA41; P2X2; P2X2a; P2RX2_HUMAN; P2RX2; ATP receptor; Purinergic receptor; purinergic receptor P2X 2; deafness, autosomal dominant 41; purinergic receptor P2X, ligand-gated ion channel, 2
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	P2RX2 (HGNC:15459)
Synonyms	P2X2

Function	ATP-gated nonselective transmembrane cation channel permeable to potassium, sodium and calcium (PubMed: 10570044 , PubMed: 31636190). Activation by extracellular ATP induces a variety of cellular responses, such as excitatory postsynaptic responses in sensory neurons, neuromuscular junctions (NMJ) formation, hearing, perception of taste and peristalsis (By similarity). In the inner ear, regulates sound transduction and auditory neurotransmission, outer hair cell electromotility, inner ear gap junctions, and K(+) recycling (PubMed: 23345450). Mediates synaptic transmission between neurons and from neurons to smooth muscle (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:P56373}. Note=Localizes to the apical membranes of hair cells in the organ of Corti.
Tissue Location	Expressed in both the central and peripheral nervous system, as well as in the pituitary gland

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.