

GSTT1 Rabbit pAb

Catalog # AP97596

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P30711
Other Accession	P30711
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27335
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human GSTT1/Glutathione S Transferase theta 1
Epitope Specificity	41-140/240
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the GST superfamily. Theta family. Contains 1 GST C-terminal domain. Contains 1 GST N-terminal domain.
SUBUNIT	Homodimer.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Glutathione S-transferase (GST) theta 1 (GSTT1) is a member of a superfamily of proteins that catalyze the conjugation of reduced glutathione to a variety of electrophilic and hydrophobic compounds. Human GSTs can be divided into five main classes: alpha, mu, pi, theta, and zeta. The theta class includes GSTT1 and GSTT2. The GSTT1 and GSTT2 share 55% amino acid sequence identity and both of them were claimed to have an important role in human carcinogenesis. The GSTT1 gene is located approximately 50kb away from the GSTT2 gene. The GSTT1 and GSTT2 genes have a similar structure, being composed of five exons with identical exon/intron boundaries. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	2952
Other Names	GSTT1_HUMAN; GSTT1; GST class-theta-1; Glutathione transferase T1-1; 2.5.1.18; glutathione S-transferase theta 1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	GSTT1
Function	Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. Acts on 1,2-epoxy-3-(4-nitrophenoxy)propane, phenethylisothiocyanate 4-nitrobenzyl chloride and 4-nitrophenethyl bromide. Displays glutathione peroxidase activity with cumene hydroperoxide.
Cellular Location	Cytoplasm.
Tissue Location	Found in erythrocyte. Expressed at low levels in liver. In lung, expressed at low levels in club cells and ciliated cells at the alveolar/bronchiolar junction. Absent from epithelial cells of larger bronchioles.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.