

MTGR1 Rabbit pAb

Catalog # AP97635

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	O43439
Other Accession	O43439
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67133
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MTGR1
Epitope Specificity	321-420/604
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the CBFA2T family.Contains 1 MYND-type zinc finger. Contains 1 TAFH (NHR1) domain.
SUBUNIT	Homooligomer and heterooligomer with MTG8. Forms a heterooligomer with the RUNX1/AML1-CBFA2T1/MTG8 chimeric protein, via an interaction with MTG8. Heterodimer with CBFA2T3.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	May function as a complex with the chimeric protein RUNX1/AML1-CBFA2T1/MTG8 which is produced in acute myeloid leukemia with the chromosomal translocation t(8;21). May thus be involved in the repression of AML1-dependent transcription and the induction of G-CSF/CSF3-dependent cell growth. May be a tumor suppressor gene candidate involved in myeloid tumors with the deletion of the 20q11 region. Tissue specificity:Ubiquitously expressed in fetal and adult tissues. Highly expressed in adult brain, heart, lung, kidney, lymph node, appendix, thymus, testis, uterus, small intestine, prostate and thymus.

Additional Information

Gene ID	9139
Other Names	EHT; MTGR1; ZMYND3; p85; A430091M07; C330013D05Rik; Cbfa2t2h; MTG8R_HUMAN; CBFA2T2; ETO homologous on chromosome 20; MTG8-like protein; MTG8-related protein 1; Myeloid translocation-related protein 1; MTG8R_MOUSE; CBFA2/RUNX1 partner transcriptional co-repressor 2; core-binding factor, runt domain, alpha subunit 2; translocated to, 2; CBFA2/RUNX1 translocation partner 2
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	CBFA2T2
Synonyms	EHT, MTGR1
Function	Transcriptional corepressor which facilitates transcriptional repression via its association with DNA-binding transcription factors and recruitment of other corepressors and histone-modifying enzymes (PubMed: 12559562 , PubMed: 15203199). Via association with PRDM14 is involved in regulation of embryonic stem cell (ESC) pluripotency (PubMed: 27281218). Involved in primordial germ cell (PCG) formation. Stabilizes PRDM14 and OCT4 on chromatin in a homooligomerization- dependent manner (By similarity). Can repress the expression of MMP7 in a ZBTB33-dependent manner (PubMed: 23251453). May function as a complex with the chimeric protein RUNX1/AML1-CBFA2T1/MTG8 (AML1-MTG8/ETO fusion protein) which is produced in acute myeloid leukemia with the chromosomal translocation t(8;21). May thus be involved in the repression of AML1-dependent transcription and the induction of G- CSF/CSF3-dependent cell growth. May be a tumor suppressor gene candidate involved in myeloid tumors with the deletion of the 20q11 region. Through heteromerization with CBFA2T3/MTG16 may be involved in regulation of the proliferation and the differentiation of erythroid progenitors by repressing the expression of TAL1 target genes (By similarity). Required for the maintenance of the secretory cell lineage in the small intestine. Can inhibit Notch signaling probably by association with RBPJ and may be involved in GFI1-mediated Paneth cell differentiation (By similarity).
Cellular Location	Nucleus.
Tissue Location	Ubiquitously expressed in fetal and adult tissues. Highly expressed in adult brain, heart, lung, kidney, lymph node, appendix, thymus, testis, uterus, small intestine, prostate and thymus

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.