

Ephrin A2 Rabbit pAb

Catalog # AP97707

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	O43921
Other Accession	O43921
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23878
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Ephrin A2/LERK6
Epitope Specificity	111-213/213
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the ephrin family. Contains 1 ephrin RBD (ephrin receptor-binding) domain.
SUBUNIT	Binds to the receptor tyrosine kinases EPHA3, EPHA4 and EPHA5. Interacts with EPHA8; activates EPHA8.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The Eph subfamily represents the largest group of receptor protein kinases identified to date. There is increasing evidence that Eph family members are involved in central nervous system function and in development. Ligands for Eph receptors include ephrin-A1 (LERK-1/B61), identified as a ligand for the EphA2 (Eck) receptor; ephrin-A2 (ELF-1), identified as a ligand for the EphA3 and EphA4 (Sek) receptors; ephrin-A3 (LERK-3), identified as a ligand for EphA5 (Ehk1) and EphA3 (Hek) receptors; ephrin-A4 (LERK-4), identified as a ligand for the EphA3 receptor; ephrin-A5 (AL-1), identified as a ligand for EphA5 (REK7); ephrin-B1 (LERK-2), identified as a ligand for the EphB1 (Elk) and EphB2 (Cek5) receptors; ephrin-B2 (LERK-5), identified as a ligand for the EphB1, EphB3 (Cek10) and EphB2 receptors; and ephrin-B3 (LERK-8), identified as a ligand for EphB1.

Additional Information

Gene ID	1943
Other Names	ELF-1; EPLG6; HEK7-L; LERK-6; LERK6; EFNA2_HUMAN; EFNA2; EPH-related receptor tyrosine kinase ligand 6 (LERK-6); HEK7 ligand (HEK7-L); ephrin A2; ephrin-A2; EPH-related receptor tyrosine kinase ligand 6
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	EFNA2
Synonyms	EPLG6, LERK6
Function	Cell surface GPI-bound ligand for Eph receptors, a family of receptor tyrosine kinases which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. Binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. With the EPHA2 receptor may play a role in bone remodeling through regulation of osteoclastogenesis and osteoblastogenesis (By similarity).
Cellular Location	Cell membrane; Lipid-anchor, GPI- anchor

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.