

DNAJB12 Rabbit pAb

Catalog # AP97728

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9NXW2
Other Accession	Q9NXW2
Reactivity	Human
Predicted	Rat, Pig, Mouse, Dog, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41860
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DNAJB12
Epitope Specificity	26-130/375
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 J domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	DNAJB12 belongs to the evolutionarily conserved DNAJ/HSP40 family of proteins, which regulate molecular chaperone activity by stimulating ATPase activity. DNAJ proteins may have up to 3 distinct domains: a conserved 70-amino acid J domain, usually at the N terminus; a glycine/phenylalanine (G/F)-rich region; and a cysteine-rich domain containing 4 motifs resembling a zinc finger domain (Ohtsuka and Hata, 2000 [PubMed 11147971]).[supplied by OMIM, Mar 2008]

Additional Information

Gene ID	54788
Other Names	DJ10; mDj10; DJB12_BOVIN; DNAJB12; DJB12_HUMAN; DJB12_MOUSE; Dnaj heat shock protein family (Hsp40) member B12; Dnaj (Hsp40) homolog, subfamily B, member 12
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	DNAJB12 {ECO:0000303 PubMed:21150129, ECO:0000312 HGNC:HGNC:14891}
Function	Acts as a co-chaperone with HSPA8/Hsc70; required to promote protein folding and trafficking, prevent aggregation of client proteins, and promote unfolded proteins to endoplasmic reticulum- associated degradation (ERAD) pathway (PubMed: 21148293 , PubMed: 21150129). Acts by determining HSPA8/Hsc70's ATPase and polypeptide-binding activities (PubMed: 21148293). Can also act independently of HSPA8/Hsc70: together with DNAJB14, acts as a chaperone that promotes maturation of potassium channels KCND2 and KCNH2 by stabilizing nascent channel subunits and assembling them into tetramers (PubMed: 27916661). While stabilization of nascent channel proteins is dependent on HSPA8/Hsc70, the process of oligomerization of channel subunits is independent of HSPA8/Hsc70 (PubMed: 27916661). When overexpressed, forms membranous structures together with DNAJB14 and HSPA8/Hsc70 within the nucleus; the role of these structures, named DJANGOs, is still unclear (PubMed: 24732912).
Cellular Location	Endoplasmic reticulum membrane; Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein. Note=Localizes to the endoplasmic reticulum membrane (PubMed:21148293, PubMed:21150129, PubMed:24732912, PubMed:27916661) When overexpressed, forms membranous structures in the nucleus (PubMed:24732912).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.