

FA83D Rabbit pAb

Catalog # AP97731

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9H4H8
Other Accession	Q9H4H8
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64424
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FA83D
Epitope Specificity	151-225/225
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the FAM83 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Representing about 2% of human DNA, chromosome 20 consists of approximately 63 million bases and 600 genes. Chromosome 20 contains a region with numerous genes expressed in the epididymis which are thought important for seminal production and some viewed as potential targets for male contraception. The PRNP gene encoding the prion protein associated with spongiform encephalopathies, like Creutzfeldt-Jakob disease, is found on chromosome 20. Amyotrophic lateral sclerosis, spinal muscular atrophy, ring chromosome 20 epilepsy syndrome and Alagille syndrome are also associated with chromosome 20. The FAM83D gene product has been provisionally designated FAM83D pending further characterization.

Additional Information

Gene ID	81610
Other Names	C20orf129; CHICA; dj616B8.3; FA83D_HUMAN; FAM83D; Spindle protein CHICA;
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SACK1D (HGNC:16122)
Function	Through the degradation of FBXW7, may act indirectly on the expression and downstream signaling of MTOR, JUN and MYC (PubMed: 24344117). May play also a role in cell proliferation through activation of the ERK1/ERK2 signaling cascade (PubMed: 25646692). May also be important for proper chromosome congression and alignment during mitosis through its interaction with KIF22 (PubMed: 18485706).
Cellular Location	Cytoplasm. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole Note=Primarily cytoplasmic during interphase, but at prophase, associates with spindle microtubules, with a clear concentration toward the spindle poles. It persists on spindle microtubules through metaphase and anaphase.
Tissue Location	Expressed in the testis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.