10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MAGEA2 Rabbit pAb

Catalog # AP97742

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	P43356
Other Accession	P43356
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35055
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MAGEA2
Epitope Specificity	21-120/314
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 MAGE domain.
SUBUNIT	Interacts with TRIM28 and UBE2H. Interacts with HDAC3. Interacts with PML (isoform PML-1, isoform PML-2, isoform PML-3, isoform PML-4 and isoform PML-5).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene is a member of the MAGEA gene family. The members of this family encode proteins with 50 to 80% sequence identity to each other. The promoters and first exons of the MAGEA genes show considerable variability, suggesting that the existence of this gene family enables the same function to be expressed under different transcriptional controls. The MAGEA genes are clustered at chromosomal location Xq28. They have been implicated in some hereditary disorders, such as dyskeratosis congenita. This gene has two identical copies at different loci. Alternatively spliced transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	266740;4101
Other Names	CT1.2; MAGE2; MAGEA2A; MAGA2_HUMAN; MAGEA2; MAGEA2B; Cancer/testis antigen 1.2 (CT1.2); MAGE-2 antigen; MAGE family member A2; melanoma antigen family A2; cancer/testis antigen family 1, member 2
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	MAGEA2
Synonyms	MAGE2, MAGEA2A
Function	Reduces p53/TP53 transactivation function through recruitment of HDAC3 to p53/TP53 transcription sites. Also represses p73/TP73 activity. Proposed to enhance ubiquitin ligase activity of RING-type zinc finger-containing E3 ubiquitin-protein ligases. In vitro enhances ubiquitin ligase activity of TRIM28 and stimulates p53/TP53 ubiquitination by TRIM28 potentially in presence of Ubl-conjugating enzyme UBE2H. Proposed to act through recruitment and/or stabilization of the Ubl-conjugating enzyme (E2) at the E3:substrate complex. May play a role in embryonal development and tumor transformation or aspects of tumor progression. In vitro promotes cell viability in melanoma cell lines. Antigen recognized on a melanoma by autologous cytolytic T-lymphocytes. Negatively regulates acetylation and sumoylation of PML and represses PML-induced p53/TP53 acetylation and activation.
Cellular Location	Nucleus. Nucleus, PML body.
Tissue Location	Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.