

Thrap3 Rabbit pAb

Catalog # AP97814

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9Y2W1
Other Accession	Q9Y2W1
Host	Rabbit
Clonality	Polyclonal
Calculated MW	108666
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Thrap3/TRAP150
Epitope Specificity	631-730/955
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBUNIT	Subunit of the large multiprotein complex TRAP.
Post-translational modifications	Phosphorylated upon DNA damage, probably by ATM or ATR.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Plays a role in transcriptional coactivation.

Additional Information

Gene ID	9967
Other Names	BCLAF2; TRAP150; 9330151F09Rik; B230333E16Rik; TR150_HUMAN; THRAP3; BCLAF1 and THRAP3 family member 2; Thyroid hormone receptor-associated protein complex 150 kDa component (Trap150); TR150_MOUSE; TR150_XENLA; thyroid hormone receptor associated protein 3; TR-associated protein, 150-kDa
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	THRAP3 (HGNC:22964)
Function	Involved in pre-mRNA splicing. Remains associated with spliced mRNA after splicing which probably involves interactions with the exon junction complex

(EJC). Can trigger mRNA decay which seems to be independent of nonsense-mediated decay involving premature stop codons (PTC) recognition. May be involved in nuclear mRNA decay. Involved in regulation of signal-induced alternative splicing. During splicing of PTPRC/CD45 is proposed to sequester phosphorylated SFPQ from PTPRC/CD45 pre-mRNA in resting T-cells. Involved in cyclin- D1/CCND1 mRNA stability probably by acting as component of the SNARP complex which associates with both the 3'end of the CCND1 gene and its mRNA. Involved in response to DNA damage. Is excluded from DNA damage sites in a manner that parallels transcription inhibition; the function may involve the SNARP complex. Initially thought to play a role in transcriptional coactivation through its association with the TRAP complex; however, it is not regarded as a stable Mediator complex subunit. Cooperatively with HELZ2, enhances the transcriptional activation mediated by PPARG, maybe through the stabilization of the PPARG binding to DNA in presence of ligand. May play a role in the terminal stage of adipocyte differentiation. Plays a role in the positive regulation of the circadian clock. Acts as a coactivator of the CLOCK-BMAL1 heterodimer and promotes its transcriptional activator activity and binding to circadian target genes (PubMed:[24043798](#)).

Cellular Location

Nucleus. Nucleus, nucleoplasm. Nucleus speckle

Tissue Location

Ubiquitous..

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.