

HAP40 Rabbit pAb

Catalog # AP97817

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P23610
Other Accession	P23610
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39103
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HAP40
Epitope Specificity	2-80/371
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene is part of a region that is repeated three times on chromosome X, once in intron 22 of the F8 gene and twice closer to the Xq telomere. This record represents the middle copy. Although its function is unknown, the observation that this gene is conserved in the mouse implies it has some function. Unlike factor VIII, this gene is transcribed abundantly in a wide variety of cell types. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	474383;474384;8263
Other Names	HAP40; DXS522E; F8A; HAP40_HUMAN; F8A1; F8A2; F8A3; CpG island protein; Factor VIII intron 22 protein; coagulation factor VIII associated 1; coagulation factor VIII-associated (intronic transcript) 1
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	F8A1
Function	RAB5A effector molecule that is involved in vesicular trafficking of early

endosomes (PubMed:[16476778](#)). Mediates the recruitment of HTT by RAB5A onto early endosomes. The HTT- F8A1/F8A2/F8A3-RAB5A complex stimulates early endosomal interaction with actin filaments and inhibits interaction with microtubules, leading to the reduction of endosome motility (PubMed:[16476778](#)).

Cellular Location

Cytoplasm. Nucleus. Early endosome. Nucleus, nuclear body {ECO:0000250|UniProtKB:Q00558}. Note=Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentrated in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity) {ECO:0000250|UniProtKB:Q00558, ECO:0000269|PubMed:16476778}

Tissue Location

Produced abundantly in a wide variety of cell types

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.