

ZHX1 Rabbit pAb

Catalog # AP97855

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9UKY1
Other Accession	Q9UKY1
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98098
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ZHX1
Epitope Specificity	551-650/873
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the ZHX family. Contains 2 C2H2-type zinc fingers. Contains 5 homeobox DNA-binding domains.
Post-translational modifications	Phosphorylated upon DNA damage, probably by ATM or ATR.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The members of the zinc fingers and homeoboxes gene family are nuclear homodimeric transcriptional repressors that interact with the A subunit of nuclear factor-Y (NF-YA) and contain two C2H2-type zinc fingers and five homeobox DNA-binding domains. This gene encodes member 1 of this gene family. In addition to forming homodimers, this protein heterodimerizes with members 2 and 3 of the zinc fingers and homeoboxes family. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the downstream chromosome 8 open reading frame 76 (C8orf76) gene. [provided by RefSeq, Feb 2011]

Additional Information

Gene ID	11244
Other Names	ZHX1_HUMAN; ZHX1; ZHX1_MOUSE; ZHX1_RAT; zinc fingers and homeoboxes 1; zinc-fingers and homeoboxes 1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ZHX1
Function	Acts as a transcriptional repressor. Increases DNMT3B- mediated repressive transcriptional activity when DNMT3B is tethered to DNA. May link molecule between DNMT3B and other co-repressor proteins.
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00108, ECO:0000269 PubMed:12237128, ECO:0000269 PubMed:17056598} Note=Colocalized in the nucleus with DNMT3B
Tissue Location	Ubiquitously expressed. Expressed in podocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.