

KIAA1543 Rabbit pAb

Catalog # AP97856

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9P1Y5
Other Accession	Q9P1Y5
Host	Rabbit
Clonality	Polyclonal
Calculated MW	134750
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KIAA1543
Epitope Specificity	851-950/1249
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the CAMSAP1 family. Contains 1 CH (calponin-homology) domain. Contains 1 CKK domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Consisting of around 63 million bases with over 1,400 genes, chromosome 19 makes up over 2% of human genomic DNA. Chromosome 19 includes a diversity of interesting genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin superfamily members including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family, and Fcα receptors. Key genes for eye color and hair color also map to chromosome 19. Peutz-Jeghers syndrome, spinocerebellar ataxia type 6, the stroke disorder CADASIL, hypercholesterolemia and insulin-dependent diabetes have been linked to chromosome 19. Translocations with chromosome 19 and chromosome 14 can be seen in some lymphoproliferative disorders and typically involve the proto-oncogene BCL3. The KIAA1543 gene product has been provisionally designated KIAA1543 pending further characterization.

Additional Information

Gene ID	57662
Other Names	KIAA1543; NEZHA; PPP1R80; CAMP3_HUMAN; CAMSAP3; Protein Nezha; calmodulin regulated spectrin associated protein family member 3; calmodulin regulated spectrin-associated protein family, member 3; protein phosphatase 1, regulatory subunit 80
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CAMSAP3 (HGNC:29307)
Function	Key microtubule-organizing protein that specifically binds the minus-end of non-centrosomal microtubules and regulates their dynamics and organization (PubMed:19041755, PubMed:23169647). Specifically recognizes growing microtubule minus-ends and autonomously decorates and stabilizes microtubule lattice formed by microtubule minus-end polymerization (PubMed:24486153). Acts on free microtubule minus-ends that are not capped by microtubule-nucleating proteins or other factors and protects microtubule minus-ends from depolymerization (PubMed:24486153). In addition, it also reduces the velocity of microtubule polymerization (PubMed:24486153). Required for the biogenesis and the maintenance of zonula adherens by anchoring the minus-end of microtubules to zonula adherens and by recruiting the kinesin KIFC3 to those junctional sites (PubMed:19041755). Required for orienting the apical-to-basal polarity of microtubules in epithelial cells: acts by tethering non-centrosomal microtubules to the apical cortex, leading to their longitudinal orientation (PubMed:26715742, PubMed:27802168). Plays a key role in early embryos, which lack centrosomes: accumulates at the microtubule bridges that connect pairs of cells and enables the formation of a non-centrosomal microtubule-organizing center that directs intracellular transport in the early embryo (By similarity). Couples non-centrosomal microtubules with actin: interaction with MACF1 at the minus ends of non-centrosomal microtubules, tethers the microtubules to actin filaments, regulating focal adhesion size and cell migration (PubMed:27693509). Plays a key role in the generation of non-centrosomal microtubules by accumulating in the pericentrosomal region and cooperating with KATNA1 to release non-centrosomal microtubules from the centrosome (PubMed:28386021). Through the microtubule cytoskeleton, also regulates the organization of cellular organelles including the Golgi and the early endosomes (PubMed:28089391). Through interaction with AKAP9, involved in translocation of Golgi vesicles in epithelial cells, where microtubules are mainly non-centrosomal (PubMed:28089391). Plays an important role in motile cilia function by facilitating proper orientation of basal bodies and formation of central microtubule pairs in motile cilia (By similarity).
Cellular Location	Cytoplasm, cytoskeleton. Cell junction, adherens junction. Cytoplasm Cytoplasm, cytoskeleton, cilium axoneme {ECO:0000250 UniProtKB:Q80VC9} Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250 UniProtKB:Q80VC9}. Note=Scattered in the cytoplasm, associated with the minus-end of microtubules and also detected at the centrosomes (PubMed:19041755, PubMed:24486153, PubMed:27693509) Decorates the minus-end of microtubules by decreasing the rate of tubulin incorporation and remaining bound (PubMed:24486153). Localizes along zonula adherens only at mature cell-cell contacts (PubMed:19041755). In early embryos, accumulates at the microtubule bridges that connect pairs of cells: this structure is present in early embryos, which lack centrosomes (By similarity). This cytokinetic bridge does not undergo stereotypical abscission after cell division (By similarity). Accumulates to the pericentrosomal region following interaction with KATNA1 (PubMed:28386021) {ECO:0000250 UniProtKB:Q80VC9, ECO:0000269 PubMed:19041755, ECO:0000269 PubMed:24486153, ECO:0000269 PubMed:27693509, ECO:0000269 PubMed:28386021}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.