

HEBP1 Rabbit pAb

Catalog # AP97862

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9NRV9
Other Accession	Q9NRV9
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21097
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HEBP1/p22HBP
Epitope Specificity	51-150/189
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the HEBP family.
SUBUNIT	Monomer.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	p22HBP, also known as HEBP1 (heme binding protein 1), HBP or HEBP, is a 189 amino acid intracellular tetrapyrrole-binding protein that assists in prevention of cellular toxicity by removing free porphyrinogens from the cell. Existing as a monomer, p22HBP localizes to cytoplasm and contains a 21 amino acid chemoattractant within its N-terminus that functions as a natural ligand for FPR3. p22HBP is a member of the HEBP family and binds N-methylprotoporphyrin and metalloporphyrins with similar affinity to porphyrinogens. The gene encoding p22HBP maps to human chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions, including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and trisomy 12p, which causes facial developmental defects and seizure disorders.

Additional Information

Gene ID	50865
Other Names	HBP; HEBP; p22HBP; HEBP1_HUMAN; HEBP1; HEBP1_MOUSE; heme binding protein 1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	HEBP1
Synonyms	HBP
Function	May bind free porphyrinogens that may be present in the cell and thus facilitate removal of these potentially toxic compound. Binds with a high affinity to one molecule of heme or porphyrins. It binds metalloporphyrins, free porphyrins and N-methylprotoporphyrin with similar affinities.
Cellular Location	Cytoplasm.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.