

STOM Rabbit pAb

Catalog # AP97864

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	P27105
Other Accession	P27105
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31731
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human STOM
Epitope Specificity	61-150/288
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the band 7/mec-2 family.
SUBUNIT	Homooligomer containing between 9 and 12 monomers. Interacts with ASIC1, ASIC2 and ASIC3. Interacts with LANCL1.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of a highly conserved family of integral membrane proteins. The encoded protein localizes to the cell membrane of red blood cells and other cell types, where it may regulate ion channels and transporters. Loss of localization of the encoded protein is associated with hereditary stomatocytosis, a form of hemolytic anemia. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2012]

Additional Information

Gene ID	2040
Other Names	BND7; EPB7; EPB72; Epb7.2; STOM_HUMAN; STOM; Erythrocyte band 7 integral membrane protein; Erythrocyte membrane protein band 7.2; Protein 7.2b; STOM_MOUSE; stomatin; erythrocyte membrane protein band 7.2 (stomatin)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	STOM (HGNC:3383)
Function	Regulates ion channel activity and transmembrane ion transport. Regulates ASIC2 and ASIC3 channel activity.
Cellular Location	Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton. Cell membrane; Lipid-anchor; Cytoplasmic side. Membrane raft. Melanosome. Cytoplasmic vesicle {ECO:0000250 UniProtKB:P54116}. Note=Localizes to juxtanuclear structure probably derived from the Golgi apparatus (PubMed:9243190) Colocalizes with cortical actin microfilaments at small plasma membrane protrusions (PubMed:9243190). Associates with alpha-granular lipid rafts (PubMed:12130500). Translocates from the alpha-granular lipid rafts to the cell membrane on thrombin activation and selectively enriched in released microvesicles (PubMed:12130500). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545).
Tissue Location	Detected in erythrocytes (at protein level). Widely expressed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.