

DTWD1 Rabbit pAb

Catalog # AP97930

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q8N5C7
Other Accession	Q8N5C7
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35248
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DTWD1
Epitope Specificity	201-304/304
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the DTW family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Encoding more than 700 genes, chromosome 15 is made up of approximately 106 million base pairs and is about 3% of the human genome. Angelman and Prader-Willi syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. In the case of Angelman syndrome, this loss is due to inactivity of the maternal 15q11-q13 encoded UBE3A gene in the brain by either chromosomal deletion or mutation. In cases of Prader-Willi syndrome, there is a partial or complete deletion of this region from the paternal copy of chromosome 15. Tay-Sachs disease is a lethal disorder associated with mutations of the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with chromosome 15 through the FBN1 gene.

Additional Information

Gene ID	56986
Other Names	MDS009; DTWD1_HUMAN; DTWD1; DTW domain-containing protein 1; 2.5.1.25; DTW motif tRNA-uridine aminocarboxypropyltransferase 1; DTW domain containing 1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	DTWD1 {ECO:0000303 PubMed:31804502, ECO:0000312 HGNC:HGNC:30926}
Function	Catalyzes the formation of 3-(3-amino-3-carboxypropyl)uridine (acp3U) at position 20 in the D-loop of several cytoplasmic tRNAs (acp3U(20)).
Cellular Location	Nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.