

SGCE Rabbit pAb

Catalog # AP97997

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	O43556
Other Accession	O43556
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49851
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SGCE
Epitope Specificity	61-160/437
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the sarcoglycan alpha/epsilon family.
DISEASE	Defects in SGCE are a cause of dystonia type 11 (DYT11) [MIM:159900]; also known as myoclonic dystonia or alcohol-responsive dystonia. DYT11 is a myoclonic dystonia. Dystonia is defined by the presence of sustained involuntary muscle contractions, often leading to abnormal postures. DYT11 is characterized by involuntary lightning jerks and dystonic movements and postures alleviated by alcohol. Inheritance is autosomal dominant. The age of onset, pattern of body involvement, presence of myoclonus and response to alcohol are all variable.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes the epsilon member of the sarcoglycan family. Sarcoglycans are transmembrane proteins that are components of the dystrophin-glycoprotein complex, which link the actin cytoskeleton to the extracellular matrix. Unlike other family members which are predominantly expressed in striated muscle, the epsilon sarcoglycan is more broadly expressed. Mutations in this gene are associated with myoclonus-dystonia syndrome. This gene is imprinted, with preferential expression from the paternal allele. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Oct 2010]

Additional Information

Gene ID	8910
Other Names	DYT11; ESG; epsilon-SG; e-SG; SGCE_BOVIN; SGCE; SGCE_HUMAN; SGCE_MOUSE; SGCE_RAT; sarcoglycan epsilon
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SGCE
Synonyms	ESG
Function	Component of the sarcoglycan complex, a subcomplex of the dystrophin-glycoprotein complex which forms a link between the F-actin cytoskeleton and the extracellular matrix.
Cellular Location	Cell membrane, sarcolemma; Single- pass membrane protein. Cytoplasm, cytoskeleton. Cell projection, dendrite. Golgi apparatus
Tissue Location	Ubiquitous.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.