

Endothelial lipase Rabbit pAb

Catalog # AP98156

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q9Y5X9
Other Accession	Q9Y5X9
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56795
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Endothelial lipase
Epitope Specificity	301-400/500
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the AB hydrolase superfamily. Lipase family.Contains 1 PLAT domain.
SUBUNIT	Head to tail homodimer.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene has substantial phospholipase activity and may be involved in lipoprotein metabolism and vascular biology. This protein is designated a member of the TG lipase family by its sequence and characteristic lid region which provides substrate specificity for enzymes of the TG lipase family. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	9388
Other Names	EDL; EL; PRO719; lipase; LIPG_HUMAN; LIPG; Endothelial cell-derived lipase (EDL EL); Phospholipase A1; 3.1.1.3; LIPG_RAT; Endothelial-derived lipase (EDL); lipase G, endothelial type; lipase, endothelial
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	LIPG
-------------	------

Function	Exerts both phospholipase and triglyceride lipase activities (PubMed: 10192396 , PubMed: 10318835 , PubMed: 12032167). More active as a phospholipase than a triglyceride lipase (PubMed: 12032167). Hydrolyzes triglycerides, both with short-chain fatty acyl groups (tributyrin) and long-chain fatty acyl groups (triolein) with similar levels of activity toward both types of substrates (PubMed: 12032167). Hydrolyzes high density lipoproteins (HDL) more efficiently than other lipoproteins (PubMed: 10192396 , PubMed: 12032167).
Cellular Location	Secreted.
Tissue Location	High level of expression in the liver, placenta, lung, thyroid, kidney, testis and in the corpus luteum of the ovary Expressed also in coronary artery endothelial cells, umbilical vein endothelial cells and in hepatocytes and osteosarcoma cell lines. Not detected in heart, brain and muscle.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.