

SEC23IP Rabbit pAb

Catalog # AP98203

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9Y6Y8
Other Accession	Q9Y6Y8
Host	Rabbit
Clonality	Polyclonal
Calculated MW	111076
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SEC23IP
Epitope Specificity	811-910/1000
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the PA-PLA1 family. Contains 1 DDHD domain. Contains 1 SAM (sterile alpha motif) domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the phosphatidic acid preferring-phospholipase A1 family. The encoded protein is localized to endoplasmic reticulum exit sites and plays a critical role in ER-Golgi transport as part of the multimeric coat protein II complex. An orthologous gene in frogs is required for normal neural crest cell development, suggesting that this gene may play a role in Waardenburg syndrome neural crest defects. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, Feb 2011]

Additional Information

Gene ID	11196
Other Names	MSTP053; P125; P125A; iPLA1A; iPLA1beta; D7Erttd373e; S23IP_HUMAN; SEC23IP; S23IP_MOUSE; SEC23 interacting protein; intracellular phospholipase A1 beta
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SEC23IP
Function	Plays a role in the organization of endoplasmic reticulum exit sites. Specifically binds to phosphatidylinositol 3-phosphate (PI(3)P), phosphatidylinositol 4-phosphate (PI(4)P) and phosphatidylinositol 5-phosphate (PI(5)P).
Cellular Location	Cytoplasmic vesicle, COPII-coated vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Endoplasmic reticulum
Tissue Location	Ubiquitously expressed with stronger levels detected in heart, liver and skeletal muscle

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.