

UNC5D Rabbit pAb

Catalog # AP98256

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q6UXZ4
Other Accession	Q6UXZ4
Host	Rabbit
Clonality	Polyclonal
Calculated MW	105880
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human UNC5H4
Epitope Specificity	112-200/953
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the unc-5 family. Contains 1 death domain. Contains 1 Ig-like (immunoglobulin-like) domain. Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 2 TSP type-1 domains. Contains 1 ZU5 domain.
SUBUNIT	Interacts with the cytoplasmic part of DCC
Post-translational modifications	Phosphorylated on cytoplasmic tyrosine residues (By similarity). Proteolytically cleaved by caspases during apoptosis. The cleavage does not take place when the receptor is associated with netrin ligand. Its cleavage by caspases is required to induce apoptosis
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The UNC5H family of proteins act as transmembrane receptors for netrin-1 and play a crucial role in axon guidance and migration of neural cells. Additionally, when cleaved by a caspase to produce an intracellular fragment containing a death domain, UNC5H receptors induce apoptosis. This activity is blocked by the binding of netrin-1. In the absence of netrin-1, UNC5H receptors act as tumor suppressors by inhibiting anchorage-independent growth and invasion, but mutation of these receptors provides a potential mechanism for tumorigenicity. The expression of UNC5H receptors is down-regulated in multiple carcinomas, including colorectal, breast, ovary, uterus, stomach, lung, and kidney cancers. UNC5H4, also known as UNC5D (unc-5 homolog D), is single-pass type I membrane protein that is a member of the UNC5H netrin receptor family. Two isoforms of UNC5H4 exist due to alternative splicing events.

Additional Information

Gene ID	137970
Other Names	PRO34692; Unc5h4; UNC5D_HUMAN; UNC5D; Protein unc-5 homolog 4; Protein unc-5 homolog D; KIAA1777; unc-5 netrin receptor D; unc-5 homolog

D (C. elegans); Netrin receptor UNC5D

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	UNC5D
Synonyms	KIAA1777, UNC5H4 {ECO:0000303 Ref.2}
Function	Receptor for the netrin NTN4 that promotes neuronal cell survival (By similarity). Plays a role in cell-cell adhesion and cell guidance. Receptor for netrin involved in cell migration. Plays a role in axon guidance by mediating axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding (By similarity). May play a role in apoptosis in response to DNA damage (PubMed: 24691657). It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (PubMed: 24519068). Mediates cell-cell adhesion via its interaction with FLRT3 on an adjacent cell (By similarity).
Cellular Location	Cell membrane; Single-pass type I membrane protein

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.