

SULT2B1 Rabbit pAb

Catalog # AP98314

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	O00204
Other Accession	O00204
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41308
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SULT2B1
Epitope Specificity	111-210/365
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the sulfotransferase 1 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Sulfotransferase enzymes catalyze the sulfate conjugation of many hormones, neurotransmitters, drugs, and xenobiotic compounds. These cytosolic enzymes are different in their tissue distributions and substrate specificities. The gene structure (number and length of exons) is similar among family members. This gene sulfates dehydroepiandrosterone but not 4-nitrophenol, a typical substrate for the phenol and estrogen sulfotransferase subfamilies. Two alternatively spliced variants that encode different isoforms have been described. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	6820
Other Names	ARCI14; HSST2; ST2B1_HUMAN; SULT2B1; Alcohol sulfotransferase; Hydroxysteroid sulfotransferase 2; Sulfotransferase family 2B member 1; Sulfotransferase family cytosolic 2B member 1 (ST2B1); 2.8.2.2; sulfotransferase family, cytosolic, 2B, member 1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SULT2B1
Synonyms	HSST2
Function	Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the sulfate conjugation. Responsible for the sulfation of cholesterol (PubMed: 12145317 , PubMed: 19589875). Catalyzes sulfation of the 3beta-hydroxyl groups of steroids, such as, pregnenolone and dehydroepiandrosterone (DHEA) (PubMed: 12145317 , PubMed: 16855051 , PubMed: 21855633 , PubMed: 9799594). Preferentially sulfonates cholesterol, while it also has significant activity with pregnenolone and DHEA (PubMed: 12145317 , PubMed: 21855633). Plays a role in epidermal cholesterol metabolism and in the regulation of epidermal proliferation and differentiation (PubMed: 28575648).
Cellular Location	Cytoplasm, cytosol. Microsome. Nucleus. Note=Phosphorylation of Ser-348 is required for translocation to the nucleus
Tissue Location	Expressed in the stratum granulosum-stratum corneum junction in the skin (at protein level) (PubMed:28575648). Expressed highly in placenta, prostate and trachea and lower expression in the small intestine and lung (PubMed:9799594)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.