

ABI1 Rabbit pAb

Catalog # AP98354

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q8IZP0
Other Accession	Q8IZP0
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55081
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ABI1
Epitope Specificity	101-200/508
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the ABI family. Contains 1 SH3 domain. Contains 1 t-SNARE coiled-coil homology domain.
SUBUNIT	Interacts with ABL1, ENAH, STX1A, SNAP25, VAMP2, EPS8, and through its N-terminus with WASF1. Part of a complex consisting of ABI1, STX1A and SNAP25. Part of a complex consisting of ABI1, EPS8 and SOS1. Interacts with SOS1, SOS2, GRB2, SPTA1 and the first SH3 domain of NCK1. Isoform 6 does not interact with NCK1. Component of the WAVE2 complex composed of ABI1, CYFIP1/SRA1, NCKAP1/NAP1 and WASF2/WAVE2.
Post-translational modifications	In vitro substrate for v-Abl. Phosphorylated on tyrosine residues after serum stimulation or induction by v-Abl.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	May act in negative regulation of cell growth and transformation by interacting with nonreceptor tyrosine kinases ABL1 and/or ABL2. In vitro, at least isoform 2 and isoform 4 suppress the transforming activity of Abelson murine leukemia virus(v-Abl) after overexpression in fibroblasts. May play a role in regulation EGF-induced Erk pathway activation. Involved in cytoskeletal reorganization and EGFR signaling. Together with EPS8 participates in transduction of signals from Ras to Rac. In vitro, a trimeric complex of ABI1, EPS8 and SOS1 exhibits Rac specific guanine nucleotide exchange factor (GEF) activity and ABI1 seems to act as an adapter in the complex. Regulates ABL1/c-Abl-mediated phosphorylation of MENA (By similarity). Recruits WASF1 to lamellipodia and there seems to regulate WASF1 protein level.

Additional Information

Gene ID	10006
Other Names	ABI-1; ABLBP4; E3B1; NAP1BP; SSH3BP; SSH3BP1; NAP1; ABI1_HUMAN; ABI1; Abelson interactor 1 (Abi-1); Abl-binding protein 4 (AbIBP4); Eps8 SH3 domain-binding protein (Eps8-binding protein); Nap1-binding protein

(Nap1BP); Spectrin SH3 domain-binding protein 1; ABI1_MOUSE; Ablphilin-1; A2VD09_RAT; ABI1_RAT; abl interactor 1; spectrin SH3 domain binding protein 1; abl-interactor 1

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ABI1 (HGNC:11320)
Synonyms	SSH3BP1
Function	May act in negative regulation of cell growth and transformation by interacting with nonreceptor tyrosine kinases ABL1 and/or ABL2. May play a role in regulation of EGF-induced Erk pathway activation. Involved in cytoskeletal reorganization and EGFR signaling. Together with EPS8 participates in transduction of signals from Ras to Rac. In vitro, a trimeric complex of ABI1, EPS8 and SOS1 exhibits Rac specific guanine nucleotide exchange factor (GEF) activity and ABI1 seems to act as an adapter in the complex. Regulates ABL1/c-Abl- mediated phosphorylation of ENAH. Recruits WASF1 to lamellipodia and there seems to regulate WASF1 protein level. In brain, seems to regulate the dendritic outgrowth and branching as well as to determine the shape and number of synaptic contacts of developing neurons.
Cellular Location	Cytoplasm. Nucleus. Cell projection, lamellipodium. Cell projection, filopodium. Cell projection, growth cone Postsynaptic density. Cytoplasm, cytoskeleton. Note=Localized to protruding lamellipodia and filopodia tips. Also localized to neuronal growth cones and synaptosomes. May shuttle from the postsynaptic densities to the nucleus (By similarity)
Tissue Location	Widely expressed, with highest expression in brain.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.