

OST alpha Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9838b

Product Information

Application	WB, FC, E
Primary Accession	Q86UW1
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB24183
Calculated MW	37735
Antigen Region	298-327

Additional Information

Gene ID	200931
Other Names	Organic solute transporter subunit alpha, OST-alpha, Solute carrier family 51 subunit alpha, SLC51A, OSTA
Target/Specificity	This OST alpha antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 298-327 amino acids from the C-terminal region of human OST alpha.
Dilution	WB~~1:1000 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	OST alpha Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SLC51A (HGNC:29955)
Synonyms	OSTA
Function	Essential component of the OST-alpha/OST-beta complex, a heterotetramer

that acts as the intestinal basolateral transporter responsible for bile acid export from enterocytes into portal blood (PubMed:[16317684](#), PubMed:[41606314](#), PubMed:[41606328](#)). The OST-alpha/OST- beta complex efficiently transports the major species of bile acids (taurocholate) (PubMed:[16317684](#), PubMed:[41606314](#), PubMed:[41606328](#)). Taurine conjugates are transported more efficiently across the basolateral membrane than glycine-conjugated bile acids (By similarity). In the complex, SLC51A/OST-alpha adopts a G protein- coupled receptor fold and constitutes a bile acid recognition site (PubMed:[41606314](#), PubMed:[41606328](#)). Can also transport steroids such as estrone 3-sulfate and dehydroepiandrosterone 3-sulfate, therefore playing a role in the enterohepatic circulation of sterols (PubMed:[16317684](#)). Able to transport eicosanoids such as prostaglandin E2 (By similarity).

Cellular Location

Basolateral cell membrane; Multi-pass membrane protein Note=Transported from the endoplasmic reticulum to the plasma membrane upon interacting with SLC51B (By similarity). Mainly restricted to the lateral and basal membranes of ileal enterocytes (By similarity)
{ECO:0000250|UniProtKB:Q8R000}


Tissue Location

Widely expressed with a high expression in ileum. Expressed in testis, colon, liver, small intestine, kidney, ovary and adrenal gland; and at low levels in heart, lung, brain, pituitary, thyroid gland, uterus, prostate, mammary gland and fat

References

Renner, O., et al. J. Lipid Res. 49(9):2045-2054(2008)
Soroka, C.J., et al. BMC Cell Biol. 9, 57 (2008)
Sun, A.Q., et al. Am. J. Physiol. Gastrointest. Liver Physiol. 292 (6), G1586-G1593 (2007)
Boyer, J.L., et al. Am. J. Physiol. Gastrointest. Liver Physiol. 290 (6), G1124-G1130 (2006)
Lim, J., et al. Cell 125(4):801-814(2006)


Images



Western blot analysis of OSTalpha Antibody (C-term) (Cat. #AP9838b) in 293 cell line lysates (35ug/lane). OST alpha (arrow) was detected using the purified Pab.

OST alpha Antibody (C-term) (Cat. #AP9838b) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

293



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.