

Zic1 Rabbit pAb

Catalog # AP98454

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q15915
Other Accession	Q15915
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48309
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Zic1
Epitope Specificity	201-286/447
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the GLI C2H2-type zinc-finger protein family. Contains 5 C2H2-type zinc fingers.
SUBUNIT	Interacts (via the C2H2-type domains 3, 4 and 5) with MDFIC (via the C2H2-type domains 3, 4 and 5). Interacts with GLI1; the interaction enhances transcription activation. Interacts with GLI2. Interacts with GLI3; the interaction enhances transcription activation
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Zic1 is a C2H2 zinc finger transcription factor that controls the expansion of neuronal precursors by inhibiting the progression of neuronal differentiation. Zic1 determines the cerebellar folial pattern by influencing proliferation in the external germinal layer (EGL). Zic1 can bind and transactivate the apolipoprotein E gene. This gene is closely linked to the gene encoding zinc finger protein of the cerebellum 4, a related family member on chromosome 3.

Additional Information

Gene ID	7545
Other Names	BAIDCS; CRS6; ZIC; ZNF201; ZIC1_HUMAN; ZIC1; Zinc finger protein 201; Zinc finger protein of the cerebellum 1; ZIC1_MOUSE; Zic family zinc finger 1; Zic family member 1; Zic family member 1 (odd-paired Drosophila homolog); Zic family member 1 (odd-paired homolog, Drosophila)
Dilution	IHC-P=1:100-500, IHC-F=1:100-500, IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ZIC1
Synonyms	ZIC, ZNF201
Function	Acts as a transcriptional activator. Involved in neurogenesis. Plays important roles in the early stage of organogenesis of the CNS, as well as during dorsal spinal cord development and maturation of the cerebellum. Involved in the spatial distribution of mossy fiber (MF) neurons within the pontine gray nucleus (PGN). Plays a role in the regulation of MF axon pathway choice. Promotes MF migration towards ipsilaterally-located cerebellar territories. May have a role in shear flow mechanotransduction in osteocytes. Retains nuclear GLI1 and GLI3 in the cytoplasm. Binds to the minimal GLI-consensus sequence 5'-TGGGTGGTC-3' (By similarity).
Cellular Location	Nucleus. Cytoplasm. Note=Localizes in the cytoplasm in presence of MDFIC overexpression.
Tissue Location	CNS. A high level expression is seen in the cerebellum. Detected in the nuclei of the cerebellar granule cell lineage from the progenitor cells of the external germinal layer to the postmigrated cells of the internal granular layer. Detected in medulloblastoma (26/29 cases), but not present in all other tumors examined.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.