

FOXN1 Rabbit pAb

Catalog # AP98495

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	O15353
Other Accession	O15353
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68925
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FOXN1
Epitope Specificity	321-420/648
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 fork-head DNA-binding domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Mutations in the winged-helix transcription factor gene at the nude locus in mice and rats produce the pleiotropic phenotype of hairlessness and athymia, resulting in a severely compromised immune system. This gene is orthologous to the mouse and rat genes and encodes a similar DNA-binding transcription factor that is thought to regulate keratin gene expression. A mutation in this gene has been correlated with T-cell immunodeficiency, the skin disorder congenital alopecia, and nail dystrophy. Alternative splicing in the 5' UTR of this gene has been observed. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	8456
Other Names	FKHL20; RONU; TIDAND; TLIND; WHN; FOXN1_HUMAN; FOXN1; Winged-helix transcription factor nude; forkhead box N1; winged-helix nude; Rowett nude
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200,Flow-Cyt=1ug/Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	FOXN1
-------------	-------

Synonyms	RONU, WHN
Function	Transcriptional regulator which regulates the development, differentiation, and function of thymic epithelial cells (TECs) both in the prenatal and postnatal thymus. Acts as a master regulator of the TECs lineage development and is required from the onset of differentiation in progenitor TECs in the developing fetus to the final differentiation steps through which TECs mature to acquire their full functionality. Regulates, either directly or indirectly the expression of a variety of genes that mediate diverse aspects of thymus development and function, including MHC Class II, DLL4, CCL25, CTSL, CD40 and PAX1. Regulates the differentiation of the immature TECs into functional cortical TECs (cTECs) and medullary TECs (mTECs). Essential for maintenance of mTECs population in the postnatal thymus. Involved in the morphogenesis and maintenance of the three-dimensional thymic microstructure which is necessary for a fully functional thymus. Plays an important role in the maintenance of hematopoiesis and particularly T lineage progenitors within the bone marrow niche with age. Essential for the vascularization of the thymus anlage. Promotes the terminal differentiation of epithelial cells in the epidermis and hair follicles, partly by negatively regulating the activity of protein kinase C (By similarity). Plays a crucial role in the early prenatal stages of T-cell ontogeny (PubMed: 21507891).
Cellular Location	Nucleus.
Tissue Location	Expressed in thymus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.