

TYW5 Rabbit pAb

Catalog # AP98505

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	A2RUC4
Other Accession	A2RUC4
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36548
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TYW5
Epitope Specificity	1-100/315
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the TYW5 family. Contains 1 JmjC domain.
SUBUNIT	Homodimer.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	C2orf60, also known as JmjC domain-containing protein C2orf60, FLJ37953 or MGC70509, is a 315 amino acid protein that exists as two alternatively spliced isoforms that are encoded by a gene that maps to human chromosome 2q33.1. As the second largest human chromosome, chromosome 2 makes up approximately 8% of the human genome and contains 237 million bases encoding over 1,400 genes. A number of genetic diseases are linked to genes on chromosome 2. Harlequin ichthyosis, a rare skin deformity, is associated with mutations in the ABCA12 gene. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alström syndrome, is related to mutations in the ALMS1 gene. Chromosome 2 contains a probable vestigial second centromere as well as vestigial telomeres, which gives credence to the hypothesis that human chromosome 2 formed as a result of an ancient fusion of two ancestral chromosomes, which are still present in modern day apes.

Additional Information

Gene ID	129450
Other Names	C2orf60; hTYW5; 1110034B05Rik; RGD1311269; TYW5_HUMAN; TYW5; tRNA yW-synthesizing protein 5; tRNA(Phe) (7-(3-amino-3-carboxypropyl)wyosine(37)-C(2))-hydroxylase; 1.14.11.42; TYW5_MOUSE; tRNA-yW synthesizing protein 5; chromosome 2 open reading frame 60
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	TYW5
Synonyms	C2orf60
Function	tRNA hydroxylase that acts as a component of the wybutosine biosynthesis pathway. Wybutosine is a hyper modified guanosine with a tricyclic base found at the 3'-position adjacent to the anticodon of eukaryotic phenylalanine tRNA. Catalyzes the hydroxylation of 7-(a- amino-a-carboxypropyl)wyosine (yW-72) into undermodified hydroxywybutosine (OHyW*). OHyW* being further transformed into hydroxywybutosine (OHyW) by LCMT2/TYW4. OHyW is a derivative of wybutosine found in higher eukaryotes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.