

GSDMD Recombinant Rabbit mAb

Catalog # AP98665

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P57764
Other Accession	P57764
Host	Rabbit
Clonality	Recombinant
Calculated MW	52801
Physical State	Liquid
Immunogen	Recombinant human GSDMD protein
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the gasdermin family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Gasdermin D is a member of the gasdermin family. Members of this family appear to play a role in regulation of epithelial proliferation. Gasdermin D has been suggested to act as a tumor suppressor. Alternatively spliced transcript variants have been described. [provided by RefSeq, Oct 2009]

Additional Information

Gene ID	79792
Other Names	DF5L; DFNA5L; FKSG10; GSDMDC1; GSDMD_HUMAN; GSDMD; Gasdermin domain-containing protein 1; gasdermin D; gasdermin domain containing 1
Dilution	WB=1:500-2000,IHC-P=1:200-800,IF=1:200-800,IHC-F=1:200-800
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	GSDMD {ECO:0000303 PubMed:26375003, ECO:0000312 HGNC:HGNC:25697}
Function	[Gasdermin-D]: Precursor of a pore-forming protein that plays a key role in host defense against pathogen infection and danger signals (PubMed: 26375003 , PubMed: 26375259 , PubMed: 27281216). This form constitutes the precursor of the pore-forming protein: upon cleavage, the released N-terminal moiety (Gasdermin-D, N-terminal) binds to membranes

and forms pores, triggering pyroptosis (PubMed:[26375003](#), PubMed:[26375259](#), PubMed:[27281216](#), PubMed:[39673710](#), PubMed:[40118151](#)).

Cellular Location

[Gasdermin-D]: Cytoplasm, cytosol. Inflammasome {ECO:0000250|UniProtKB:Q9D8T2}. Note=In response to a canonical inflammasome stimulus, such as nigericin, recruited to NLRP3 inflammasome with similar kinetics to that of uncleaved CASP1 precursor. {ECO:0000250|UniProtKB:Q9D8T2} [Gasdermin-D, N-terminal]: Cytoplasm, cytosol. Note=(Microbial infection) Upon infection by M.tuberculosis, localization to cell membrane is prevented by M.tuberculosis phosphatase PtpB that catalyzes dephosphorylation of phosphatidylinositol (4,5)-bisphosphate and phosphatidylinositol 4- phosphate, thereby inhibiting the membrane targeting of Gasdermin-D, N- terminal and subsequent cytokine release and pyroptosis [Gasdermin-D, C-terminal]: Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9D8T2}

Tissue Location

Expressed in the suprabasal cells of esophagus, as well as in the isthmus/neck, pit, and gland of the stomach, suggesting preferential expression in differentiating cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.