

SP7 | Osterix Recombinant Rabbit mAb

Catalog # AP98666

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q8TDD2
Other Accession	Q8TDD2
Host	Rabbit
Clonality	Recombinant
Calculated MW	44994
Physical State	Liquid
Immunogen	Recombinant human Sp7 protein
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the Sp1 C2H2-type zinc-finger protein family. Contains 3 C2H2-type zinc fingers.
SUBUNIT	Interacts with NO66; the interaction is direct and inhibits transcription activator activity.
DISEASE	Osteogenesis imperfecta 12 (OI12) [MIM:613849]: A form of osteogenesis imperfecta, a connective tissue disorder characterized by low bone mass, bone fragility and susceptibility to fractures after minimal trauma. Disease severity ranges from very mild forms without fractures to intrauterine fractures and perinatal lethality. Extraskeletal manifestations, which affect a variable number of patients, are dentinogenesis imperfecta, hearing loss, and blue sclerae. OI12 is an autosomal recessive form characterized by recurrent fractures, mild bone deformations, generalized osteoporosis, delayed teeth eruption, no dentinogenesis imperfecta, normal hearing, and white sclerae. Note=The disease is caused by mutations affecting the gene represented in this entry.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the Sp subfamily of Sp/XKLF transcription factors. Sp family proteins are sequence-specific DNA-binding proteins characterized by an amino-terminal trans-activation domain and three carboxy-terminal zinc finger motifs. This protein is a bone specific transcription factor and is required for osteoblast differentiation and bone formation.[provided by RefSeq, Jul 2010]

Additional Information

Gene ID	121340
Other Names	OI11; OI12; OSX; osterix; 6430578P22Rik; C22; SP7_HUMAN; SP7; Zinc finger protein osterix; Q5RM08_MOUSE; SP7_MOUSE; Q6IMK1_RAT; Sp7 transcription factor

Dilution	WB=1:500-1000,IHC-P=1:200-1000,IF=1:200-1000,IHC-F=1:200-1000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SP7
Synonyms	OSX
Function	Transcriptional activator essential for osteoblast differentiation (PubMed: 23457570). Binds to SP1 and EKLF consensus sequences and to other G/C-rich sequences (By similarity).
Cellular Location	Nucleus {ECO:0000250 UniProtKB:Q8VI67}.
Tissue Location	Restricted to bone-derived cell.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.