10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Angiopoietin 1 Recombinant Rabbit mAb

Catalog # AP98693

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q15389
Other Accession	Q15389
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Recombinant
Calculated MW	57513
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 fibrinogen C-terminal domain.
SUBUNIT	Homooligomer. Interacts with TEK/TIE2.
Post-translational modifications	Glycosylated.
Background Descriptions	function:Binds and activates TIE2 receptor by inducing its tyrosine phosphorylation. Implicated in endothelial developmental processes later and distinct from that of VEGF. Appears to play a crucial role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme. Mediates blood vessel maturation/stability. It may play an important role in the heart early development.,miscellaneous:It may have a potential therapeutic utility since it can be used for specifically targeting tumor vasculature or for promoting angiogenic processes in certain organs such as an ischemic heart.,online information:Angiopoietin entry,PTM:Glycosylated.,similarity:Contains 1 fibrinogen C-terminal domain.,

Additional Information

Gene ID	284
Other Names	AGP1; AGPT; AGPT-1; ANG1; HAE5; 1110046O21Rik; Ang-1; ANGP1_HUMAN; ANGPT1; KIAA0003; ANGP1_MOUSE; angiopoietin 1; Angiopoietin-1
Dilution	IHC-P=1:50-200,IHC-F=1:50-200,WB=1:500-2000,IF=1:50-200,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ANGPT1
Synonyms	KIAA0003
Function	Binds and activates TEK/TIE2 receptor by inducing its dimerization and tyrosine phosphorylation. Plays an important role in the regulation of angiogenesis, endothelial cell survival, proliferation, migration, adhesion and cell spreading, reorganization of the actin cytoskeleton, but also maintenance of vascular quiescence. Required for normal angiogenesis and heart development during embryogenesis. After birth, activates or inhibits angiogenesis, depending on the context. Inhibits angiogenesis and promotes vascular stability in quiescent vessels, where endothelial cells have tight contacts. In quiescent vessels, ANGPT1 oligomers recruit TEK to cell- cell contacts, forming complexes with TEK molecules from adjoining cells, and this leads to preferential activation of phosphatidylinositol 3-kinase and the AKT1 signaling cascades. In migrating endothelial cells that lack cell-cell adhesions, ANGT1 recruits TEK to contacts with the extracellular matrix, leading to the formation of focal adhesion complexes, activation of PTK2/FAK and of the downstream kinases MAPK1/ERK2 and MAPK3/ERK1, and ultimately to the stimulation of sprouting angiogenesis. Mediates blood vessel maturation/stability. Implicated in endothelial developmental processes later and distinct from that of VEGF. Appears to play a crucial role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.