

SAHH Rabbit pAb

Catalog # AP98712

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P23526
Other Accession	P23526
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47716
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SAHH
Epitope Specificity	25-125/432
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the adenosylhomocysteinase family.
SUBUNIT	Homotetramer.
DISEASE	Defects in AHCY are the cause of hypermethioninemia with S-adenosylhomocysteine hydrolase deficiency (HMAHCHD) [MIM:613752]. A metabolic disorder characterized by hypermethioninemia associated with failure to thrive, mental and motor retardation, facial dysmorphism with abnormal hair and teeth, and myocardiopathy.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Adenosylhomocysteine is a competitive inhibitor of S-adenosyl-L-methionine-dependent methyl transferase reactions; therefore adenosylhomocysteinase may play a key role in the control of methylations via regulation of the intracellular concentration of adenosylhomocysteine.

Additional Information

Gene ID	191
Other Names	SAHH; adoHcyase; CuBP; SAHH_HUMAN; AHCY; S-adenosyl-L-homocysteine hydrolase; 3.13.2.1; SAHH_MOUSE; Liver copper-binding protein; SAHH_RAT; adenosylhomocysteinase; S-adenosylhomocysteine hydrolase
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1ug /Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	AHCY
Synonyms	SAHH
Function	Catalyzes the hydrolysis of S-adenosyl-L-homocysteine to form adenosine and homocysteine (PubMed: 10933798). Binds copper ions (By similarity).
Cellular Location	Cytoplasm. Melanosome. Nucleus. Endoplasmic reticulum. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.