

CHST12 Rabbit pAb

Catalog # AP98755

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9NRB3
Other Accession	Q9NRB3
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48414
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CHST12
Epitope Specificity	201-300/414
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the sulfotransferase 2 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene belongs to the sulfotransferase 2 family. It is localized to the golgi membrane, and catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine (GalNAc) residue of chondroitin and desulfated dermatan sulfate. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage, and is distributed on the surfaces of many cells and extracellular matrices. Alternatively spliced transcript variants differing only in their 5' UTRs have been found for this gene. [provided by RefSeq, Aug 2011]

Additional Information

Gene ID	55501
Other Names	C4S-2; C4ST-2; C4ST2; CHSTC_HUMAN; CHST12; Chondroitin 4-O-sulfotransferase 2; Chondroitin 4-sulfotransferase 2 (C4ST-2 C4ST2); Sulfotransferase Hlo; 2.8.2.5; carbohydrate sulfotransferase 12; carbohydrate (chondroitin 4) sulfotransferase 12; chondroitin 4-sulfotransferase 2
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CHST12
Function	Catalyzes the transfer of sulfate to position 4 of the N- acetylgalactosamine (GalNAc) residue of chondroitin and desulfated dermatan sulfate. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. Activity toward partially desulfated dermatan sulfate is however lower. Does not form 4, 6-di-O- sulfated GalNAc when chondroitin sulfate C is used as an acceptor.
Cellular Location	Golgi apparatus membrane; Single- pass type II membrane protein
Tissue Location	Widely expressed. Expressed a high level in spinal chord, heart, spleen, thyroid, pituitary gland, adrenal gland, peripheral blood leukocytes, thymus, lung, small intestine, fetal kidney, fetal spleen and fetal lung.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.