

AIM2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9876a

Product Information

Application	WB, FC, IF, E
Primary Accession	O14862
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB23659
Calculated MW	38954
Antigen Region	4-32

Additional Information

Gene ID	9447
Other Names	Interferon-inducible protein AIM2, Absent in melanoma 2, AIM2
Target/Specificity	This AIM2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 4-32 amino acids from the N-terminal region of human AIM2.
Dilution	WB~~1:1000 FC~~1:10~50 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	AIM2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	AIM2 {ECO:0000303 PubMed:9242382, ECO:0000312 HGNC:HGNC:357}
Function	Sensor component of the AIM2 inflammasome, which mediates inflammasome activation in response to the presence of double-stranded DNA (dsDNA) in the cytosol, leading to subsequent pyroptosis (PubMed: 17726700 , PubMed: 19158675 , PubMed: 19158676 ,

PubMed:[19158679](#), PubMed:[20566831](#), PubMed:[23530044](#), PubMed:[26197926](#), PubMed:[26583071](#), PubMed:[29440442](#), PubMed:[33980849](#), PubMed:[37364111](#)). Inflammasomes are supramolecular complexes that assemble in the cytosol in response to pathogens and other damage-associated signals and play critical roles in innate immunity and inflammation (PubMed:[17726700](#), PubMed:[19158675](#), PubMed:[19158676](#), PubMed:[19158679](#), PubMed:[20566831](#), PubMed:[26197926](#), PubMed:[29440442](#), PubMed:[33980849](#)). Acts as a recognition receptor (PRR): specifically recognizes and binds dsDNA in the cytosol, and mediates the formation of the inflammasome polymeric complex composed of AIM2, CASP1 and PYCARD/ASC (PubMed:[17726700](#), PubMed:[19158675](#), PubMed:[19158676](#), PubMed:[19158679](#), PubMed:[20566831](#), PubMed:[26197926](#), PubMed:[29440442](#), PubMed:[33980849](#)). Recruitment of pro-caspase-1 (proCASP1) to the AIM2 inflammasome promotes caspase-1 (CASP1) activation, which subsequently cleaves and activates inflammatory cytokines IL1B and IL18 and gasdermin-D (GSDMD), promoting cytokine secretion (PubMed:[17726700](#), PubMed:[19158675](#), PubMed:[19158676](#), PubMed:[19158679](#), PubMed:[20566831](#)). In some cells, CASP1 activation mediates cleavage and activation of GSDMD, triggering pyroptosis without promoting cytokine secretion (PubMed:[19158675](#), PubMed:[19158676](#)). Detects cytosolic dsDNA of viral and bacterial origin in a non-sequence-specific manner (PubMed:[17726700](#), PubMed:[19158675](#), PubMed:[19158676](#), PubMed:[19158679](#), PubMed:[20566831](#), PubMed:[26197926](#), PubMed:[26583071](#), PubMed:[29440442](#), PubMed:[33980849](#)). Involved in the DNA damage response caused by acute ionizing radiation by mediating pyroptosis of intestinal epithelial cells and bone marrow cells in response to double-strand DNA breaks (By similarity). Mechanistically, AIM2 senses DNA damage in the nucleus to mediate inflammasome assembly and inflammatory cell death (By similarity). Also acts as a regulator of neurodevelopment via its role in the DNA damage response: acts by promoting neural cell death in response to DNA damage in the developing brain, thereby purging genetically compromised cells of the central nervous system (By similarity). Pyroptosis mediated by the AIM2 inflammasome in response to DNA damage is dependent on GSDMD without involving IL1B and IL18 cytokine secretion (By similarity). Also acts as a mediator of pyroptosis, necroptosis and apoptosis (PANoptosis), an integral part of host defense against pathogens, in response to bacterial infection (By similarity). Can also trigger PYCARD/ASC- dependent, caspase-1-independent cell death that involves caspase-8 (CASP8) (By similarity).

Cellular Location

Cytoplasm. Inflammasome. Nucleus. Note=Activated inflammasomes can aggregate in the cytosol as speck-like particles (PubMed:[19158675](#), PubMed:[19158676](#), PubMed:[19158679](#)). Activated inflammasomes can also aggregate in the nucleus in response to DNA damage: AIM2 is recruited to double-strand DNA breaks and mediates activation of the AIM2 inflammasome (By similarity). {ECO:0000250|UniProtKB:Q91VJ1, ECO:0000269|PubMed:[19158675](#), ECO:0000269|PubMed:[19158676](#), ECO:0000269|PubMed:[19158679](#)}

Tissue Location

Expressed in spleen, small intestine, peripheral blood leukocytes, and testis.


Background

AIM2 is a member of the IFI202X /IFI16 family. It plays a putative role in tumorigenic reversion and may control cell proliferation. Interferon-gamma induces expression of AIM2.

References

Patsos, G., et al. Int. J. Cancer 126(8):1838-1849(2010)
Patsos, G., et al. Glycobiology 19(7):726-734(2009)
Fernandes-Alnemri, T., et al. Nature 458(7237):509-513(2009)
Hornung, V., et al. Nature 458(7237):514-518(2009)
Woerner, S.M., et al. Genes Chromosomes Cancer 46(12):1080-1089(2007)
Cresswell, K.S., et al. Biochem. Biophys. Res. Commun. 326(2):417-424(2005)

Images



All lanes: Anti-AIM2 Antibody (N-term) at 1:2000 dilution + HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 39 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.