

KLHL20 Rabbit pAb

Catalog # AP98785

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9Y2M5
Other Accession	Q9Y2M5
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67955
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KLHL20
Epitope Specificity	13-100/609
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 BACK (BTB/Kelch associated) domain. Contains 1 BTB (POZ) domain. Contains 6 Kelch repeats.
SUBUNIT	Component of the BCR(KLHL20) E3 ubiquitin ligase complex, at least composed of CUL3, KLHL20 and RBX1. Interacts with DAPK1. Interacts with F-actin. Interacts with IFN-gamma (IFNG).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex involved in interferon response. The BCR(KLHL20) E3 ubiquitin ligase complex mediates the ubiquitination of DAPK1, leading to its degradation by the proteasome, thereby acting as a negative regulator of apoptosis. Also acts as a regulator of endothelial migration during angiogenesis by controlling the activation of Rho GTPases.

Additional Information

Gene ID	27252
Other Names	KLH20_HUMAN; KLHL20; Kelch-like ECT2-interacting protein; Kelch-like protein X; KLEIP; KLHLX; KHLHX; kelch like family member 20; kelch-like 20 (Drosophila)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	KLHL20 (HGNC:25056)
Function	Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex involved in interferon response and anterograde Golgi to endosome transport. The BCR(KLHL20) E3 ubiquitin ligase complex mediates the ubiquitination of DAPK1, leading to its degradation by the proteasome, thereby acting as a negative regulator of apoptosis (PubMed: 20389280). The BCR(KLHL20) E3 ubiquitin ligase complex also specifically mediates 'Lys-33'-linked ubiquitination (PubMed: 24768539). Involved in anterograde Golgi to endosome transport by mediating 'Lys-33'-linked ubiquitination of CORO7, promoting interaction between CORO7 and EPS15, thereby facilitating actin polymerization and post-Golgi trafficking (PubMed: 24768539). Also acts as a regulator of endothelial migration during angiogenesis by controlling the activation of Rho GTPases. The BCR(KLHL20) E3 ubiquitin ligase complex acts as a regulator of neurite outgrowth by mediating ubiquitination and degradation of PDZ-RhoGEF/ARHGEF11 (PubMed: 21670212). In case of tumor, the BCR(KLHL20) E3 ubiquitin ligase complex is involved in tumor hypoxia: following hypoxia, the BCR(KLHL20) complex mediates ubiquitination and degradation of PML, potentiating HIF-1 signaling and cancer progression (PubMed: 21840486).
Cellular Location	Cytoplasm, perinuclear region. Golgi apparatus, trans-Golgi network. Cell projection, axon {ECO:0000250 UniProtKB:D3Z8N4}. Cell projection, dendrite {ECO:0000250 UniProtKB:D3Z8N4}. Note=Localizes in the perinuclear region in normal conditions. Following IFN-alpha or IFN-gamma treatment, it is relocalized and sequestered to the PML nuclear bodies, preventing DAPK1 ubiquitination (PubMed:20389280)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.