

GALNT9 Rabbit pAb

Catalog # AP98810

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9HCQ5
Other Accession	Q9HCQ5
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68359
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human GALNT9
Epitope Specificity	401-500/603
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the glycosyltransferase 2 family. GalNAc-T subfamily. Contains 1 ricin B-type lectin domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GalNAc-T) family of enzymes. GalNAc-Ts initiate mucin-type O-linked glycosylation in the Golgi apparatus by catalyzing the transfer of GalNAc to serine and threonine residues on target proteins. They are characterized by an N-terminal transmembrane domain, a stem region, a luminal catalytic domain containing a GT1 motif and Gal/GalNAc transferase motif, and a C-terminal ricin/lectin-like domain. GalNAc-Ts have different, but overlapping, substrate specificities and patterns of expression. This gene is expressed specifically in the brain, with highest expression in the cerebellum. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	50614
Other Names	GALNAC-T9; GALNACT9; WBSCR17; si:ch211-126h20.1; GALT9_HUMAN; GALNT9; Polypeptide GalNAc transferase 9 (GalNAc-T9 pp-GaNTase 9); Protein-UDP acetylgalactosaminyltransferase 9; UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase 9; 2.4.1.41; polypeptide N-acetylgalactosaminyltransferase 9; UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 9 (GalNAc-T9); polypeptide GalNAc transferase 9
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	GALNT9
Function	Catalyzes the initial reaction in O-linked oligosaccharide biosynthesis, the transfer of an N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor. Does not glycosylate apomucin or SDC3.
Cellular Location	Golgi apparatus membrane; Single- pass type II membrane protein
Tissue Location	Specifically expressed in brain. Not expressed in heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon and leukocyte. In brain, it is expressed in cerebellum, frontal lobe, temporal lobe, putamen and spinal cord, weakly expressed in cerebral cortex. Not expressed in medulla and occipital pole

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.