

NDUAB Rabbit pAb

Catalog # AP98832

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q86Y39
Other Accession	Q86Y39
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14852
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NDUAB
Epitope Specificity	31-130/141
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the complex I NDUFA11 subunit family.
DISEASE	Defects in NDUFA11 are a cause of mitochondrial complex I deficiency (MT-C1D) [MIM:252010]. A disorder of the mitochondrial respiratory chain that causes a wide range of clinical manifestations, from lethal neonatal disease to adult-onset neurodegenerative disorders. Phenotypes include macrocephaly with progressive leukodystrophy, non-specific encephalopathy, cardiomyopathy, myopathy, liver disease, Leigh syndrome, Leber hereditary optic neuropathy and some forms of Parkinson disease.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a subunit of the membrane-bound mitochondrial complex I. Complex I is composed of numerous subunits and functions as the NADH-ubiquinol reductase of the mitochondrial electron transport chain. Mutations in this gene are associated with severe mitochondrial complex I deficiency. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Oct 2010]

Additional Information

Gene ID	126328
Other Names	B14.7; CI-B14.7; MC1DN14; 2010012C24Rik; NDUAB_HUMAN; NDUFA11; Complex I-B14.7 (CI-B14.7); NADH-ubiquinone oxidoreductase subunit B14.7; NDUAB_MOUSE; NADH:ubiquinone oxidoreductase subunit A11; NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 11, 14.7kDa; complex I B14.7 subunit
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	NDUFA11
Function	Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.
Cellular Location	Mitochondrion inner membrane; Multi-pass membrane protein; Matrix side

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.