

Synapsin III Rabbit pAb

Catalog # AP98893

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	O14994
Other Accession	O14994
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63303
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Synapsin III
Epitope Specificity	251-350/580
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the synapsin family.
SUBUNIT	Interacts with CAPON
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Synapsins are synaptic vesicle-associated phosphoproteins that regulate synaptic vesicle exocytosis and may be involved in synaptogenesis. Evidence suggests that Synapsin I, Synapsin II and Synapsin IIIa are ATP-binding proteins that are regulated by Ca ²⁺ and calmodulin binding. Ca ²⁺ has been shown to stimulate ATP binding to Synapsin I, to have no effect on Synapsin II and to inhibit Synapsin III. Synapsin I and Synapsin II both undergo alternative splicing to produce two forms of each protein, Synapsin Ia and Ib and Synapsin IIa and IIb, respectively. Synapsin III gives rise to at least three isoforms: Synapsin IIIa, IIIb and IIIc. Synapsin III plays unique roles both in early axon outgrowth and in the regulation of synaptic vesicle trafficking. In cultured mouse hippocampal neurons, synapsin III is expressed early during development, with levels peaking seven days after plating and declining thereafter. Synapsin III is highly concentrated in growth cones.

Additional Information

Gene ID	8224
Other Names	SYN3_HUMAN; SYN3; Synapsin III; SYN3_RAT; Synapsin-3
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SYN3
Function	May be involved in the regulation of neurotransmitter release and synaptogenesis.
Cellular Location	Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane; Peripheral membrane protein; Cytoplasmic side Note=Peripheral membrane protein localized to the cytoplasmic surface of synaptic vesicles
Tissue Location	Neuron specific. Detected predominantly in brain.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.