

TRIP Rabbit pAb

Catalog # AP98946

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9BWF2
Other Accession	Q9BWF2
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53294
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TRIP
Epitope Specificity	201-300/469
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 RING-type zinc finger.
SUBUNIT	Binds TRAF1 and TRAF2 and is part of the receptor-TRAF signaling complex. Interacts with CYLD.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	RING finger motif and a putative coiled-coil domain. A similar murine protein interacts with TNFR-associated factor 1 (TRAF1), TNFR-associated factor 2 (TRAF2), and cylindromatosis. The interaction with TRAF2 inhibits TRAF2-mediated nuclear factor kappa-B, subunit 1 activation that is required for cell activation and protection against apoptosis.

Additional Information

Gene ID	10293
Other Names	RNF206; SCKL9; TRIP; TRAFIP_HUMAN; TRAFIP; RING finger protein 206; TRAF-interacting protein; 2.3.2.27; TRAF interacting protein
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TRAFIP {ECO:0000303 PubMed:26595769, ECO:0000312 HGNC:HGNC:30764}
-------------	---

Function

E3 ubiquitin ligase required to protect genome stability in response to replication stress (PubMed:[25335891](#), PubMed:[26595769](#), PubMed:[26711499](#), PubMed:[26781088](#), PubMed:[27462463](#), PubMed:[31545170](#)). Acts as a key regulator of interstrand cross-link repair, which takes place when both strands of duplex DNA are covalently tethered together, thereby blocking replication and transcription (By similarity). Controls the choice between the two pathways of replication-coupled interstrand-cross-link repair by mediating ubiquitination of MCM7 subunit of the CMG helicase complex (By similarity). Short ubiquitin chains on MCM7 promote recruitment of DNA glycosylase NEIL3 (By similarity). If the interstrand cross-link cannot be cleaved by NEIL3, the ubiquitin chains continue to grow on MCM7, promoting the unloading of the CMG helicase complex by the VCP/p97 ATPase, enabling the Fanconi anemia DNA repair pathway (By similarity). Only catalyzes ubiquitination of MCM7 when forks converge (By similarity). Also involved in the repair of covalent DNA-protein cross-links (DPCs) during DNA synthesis: promotes ubiquitination of DPCs, leading to their degradation by the proteasome (By similarity). Has also been proposed to play a role in promoting translesion synthesis by mediating the assembly of 'Lys-63'-linked poly-ubiquitin chains on the Y-family polymerase POLN in order to facilitate bypass of DNA lesions and preserve genomic integrity (PubMed:[24553286](#)). The function in translesion synthesis is however controversial (PubMed:[26595769](#)). Acts as a regulator of the spindle assembly checkpoint (PubMed:[25335891](#)). Also acts as a negative regulator of innate immune signaling by inhibiting activation of NF-kappa-B mediated by TNF (PubMed:[22945920](#)). Negatively regulates TLR3/4- and RIG-I-mediated IRF3 activation and subsequent IFNB1 production and cellular antiviral response by promoting 'Lys-48'-linked polyubiquitination of TNK1 leading to its proteasomal degradation (PubMed:[22945920](#)).

Cellular Location

Nucleus, nucleoplasm. Nucleus, nucleolus. Chromosome. Cytoplasm
Cytoplasm, perinuclear region. Note=In the nucleus, found in close proximity to PCNA, suggesting localization at replication foci (PubMed:[26595769](#)). Localizes to DNA damage sites in response to replication stress (PubMed:[26595769](#), PubMed:[26711499](#), PubMed:[26781088](#)).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.