

TFPI-2 Rabbit pAb

Catalog # AP98979

Product Information

Application	IHC-P, IHC-F, IF
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25 kDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from mouse TFPI-2
Epitope Specificity	1-100/230
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 3 BPTI/Kunitz inhibitor domains.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	TFPI2 may play a role in the regulation of plasmin-mediated matrix remodeling. It inhibits trypsin, plasmin, factor VIIa/tissue factor and weakly factor Xa, but it has no effect on thrombin. TFPI2 is specific to umbilical vein endothelial cells, liver, placenta, heart, pancreas, and maternal serum at advanced pregnancy. This inhibitor contains three inhibitory domains. TFPI2 contains 3 BPTI/Kunitz inhibitor domains.

Additional Information

Other Names	PP5; REF1; TFPI-2; TFPI2_HUMAN; TFPI2; Placental protein 5 (PP5); tissue factor pathway inhibitor 2
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Name	TFPI-2 Rabbit pAb
Function	May play a role in the regulation of plasmin-mediated matrix remodeling. Inhibits trypsin, plasmin, factor VIIa/tissue factor and weakly factor Xa. Has no effect on thrombin.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.