

IL17E Rabbit pAb

Catalog # AP98980

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9H293
Other Accession	Q9H293
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20330
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human IL17E
Epitope Specificity	431-434/644
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the IL-17 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a cytokine that shares sequence similarity with interleukin 17. This cytokine can induce NF-kappaB activation, and stimulate the production of interleukin8. Both this cytokine and interleukin 17B are ligands for the cytokine receptor IL17BR. Studies of a similar gene in mice suggest that this cytokine may be a pro-inflammatory cytokine favoring the Th2-type immune response. Alternative splicing results in multiple transcript variants. [provided by RefSeq].

Additional Information

Gene ID	64806
Other Names	IL17E; IL-17e; IL-25; IL25_HUMAN; IL25; Interleukin-17E (IL-17E); interleukin 25; interleukin 17E
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	IL25
Synonyms	IL17E

Function	Cytokine produced by various cells such as eosinophils, T- helper type 2 (Th2) cells or epithelial cells that plays a role in internal safety of adaptive immune responses by regulating cytokine production (PubMed: 15860795 , PubMed: 25821217). Promotes and augments T- helper type 2 responses locally or systemically (PubMed: 25821217). Exerts its activity via its receptor composed of IL17RA and IL17RB for signal transduction (By similarity). In turn, stimulates the JAK2- STAT5A pathway and promotes the secretion of type-2 associated cytokines including IL4, IL9 and IL13 (PubMed: 25821217). Also induces the release of IL8, and IL6 from eosinophils through the combined activation of MAPK and NF-kappa-B pathways (PubMed: 15860795). Inhibits the differentiation of T-helper (Th17) cells via the production of IL4, IL5 and IL13 (PubMed: 11754819).
Cellular Location	Secreted.
Tissue Location	Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and trachea

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.