

Sprouty 4 Rabbit pAb

Catalog # AP99025

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9C004
Other Accession	Q9C004
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32541
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Sprouty 4
Epitope Specificity	1-100/299
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the sprouty family. Contains 1 SPR (sprouty) domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	SPRY4 is an inhibitor of the receptor-transduced mitogen-activated protein kinase (MAPK) signaling pathway. It is positioned upstream of RAS (see HRAS; MIM 190020) activation and impairs the formation of active GTP-RAS (Leeksma et al., 2002 [PubMed 12027893]).[supplied by OMIM, Mar 2008]

Additional Information

Gene ID	81848
Other Names	HH17; A030006O18Rik; sprouty4; SPY4_BOVIN; SPRY4; Spry-4; SPY4_HUMAN; SPY4_MOUSE; sprouty RTK signaling antagonist 4; sprouty homolog 4 (Drosophila)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SPRY4
Function	Suppresses the insulin receptor and EGFR-transduced MAPK signaling pathway, but does not inhibit MAPK activation by a constitutively active

mutant Ras (PubMed:[12027893](#)). Probably impairs the formation of GTP-Ras (PubMed:[12027893](#)). Inhibits Ras-independent, but not Ras-dependent, activation of RAF1 (PubMed:[12717443](#)). Represses integrin-mediated cell spreading via inhibition of TESK1-mediated phosphorylation of cofilin (PubMed:[15584898](#)).

Cellular Location

Cytoplasm. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Note=Found in the cytoplasm in unstimulated cells but is translocated to the membrane ruffles in cells stimulated with EGF (epidermal growth factor) (By similarity). Colocalizes with TESK1 in vesicular spots in the cytoplasm (PubMed:15584898) {ECO:0000250|UniProtKB:Q9WTP2, ECO:0000269|PubMed:15584898}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.