10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

UNC5C Rabbit pAb

Catalog # AP99031

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	O95185
Other Accession	O95185
Predicted	Rat, Pig, Human, Mouse, Rabbit, Chicken, Dog, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	103146
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human UNC5C
Epitope Specificity	201-288/931
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the unc-5 family. Contains 1 death domain. Contains 1 Ig-like (immunoglobulin-like) domain. Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 2 TSP type-1 domains. Contains 1 ZU5 domain.
SUBUNIT	Interacts with the cytoplasmic part of DCC
Post-translational modifications	Phosphorylated on different cytoplasmic tyrosine residues. Phosphorylation of Tyr-568 leads to an interaction with PTPN11 phosphatase, suggesting that its activity is regulated by phosphorylation/dephosphorylation. Tyrosine phosphorylation is netrin-dependent (By similarity). Proteolytically cleaved by caspases during apoptosis. The cleavage does not take place when the receptor is associated with netrin ligand. Its cleavage by caspases is required to induce apoptosis
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene product belongs to the UNC-5 family of netrin receptors. Netrins are secreted proteins that direct axon extension and cell migration during neural development. They are bifunctional proteins that act as attractants for some cell types and as repellents for others, and these opposite actions are thought to be mediated by two classes of receptors. The UNC-5 family of receptors mediate the repellent response to netrin; they are transmembrane proteins containing 2 immunoglobulin (Ig)-like domains and 2 type I thrombospondin motifs in the extracellular region. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	8633
Other Names	UNC5H3; B130051O18Rik; rcm; UNC5C_HUMAN; UNC5C; Protein unc-5 homolog 3; Protein unc-5 homolog C; UNC5C_MOUSE; Rostral cerebellar

	malformation protein; UNC5C_RAT; unc-5 netrin receptor C; unc5 (C.elegans homolog) c; unc-5 homolog C (C. elegans)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	UNC5C
Synonyms	UNC5H3
Function	Receptor for netrin required for axon guidance (By similarity). Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding (By similarity). NTN1/Netrin-1 binding might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed:28483977). Might also collaborate with DSCAM in NTN1-mediated axon repulsion independently of DCC (By similarity). Also involved in corticospinal tract axon guidance independently of DCC (By similarity). Involved in dorsal root ganglion axon projection towards the spinal cord (PubMed:28483977). It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (By similarity).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Cell surface. Synapse, synaptosome {ECO:0000250|UniProtKB:Q761X5}. Cell projection, axon {ECO:0000250|UniProtKB:O08747}. Cell projection, dendrite {ECO:0000250|UniProtKB:O08747}. Cell projection, growth cone {ECO:0000250|UniProtKB:O08747}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:O08747}. Cell projection, filopodium {ECO:0000250|UniProtKB:O08747}
Tissue Location	Mainly expressed in brain (PubMed:9782087). Expressed in temporal lobe cortical neurons and in neurons of the hippocampal pyramidal layer (PubMed:25419706). Also expressed in kidney (PubMed:9782087). Not expressed in developing or adult lung (PubMed:9782087).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.