

PCTAIRE2 Rabbit pAb

Catalog # AP99077

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q00537
Other Accession	Q00537
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59582
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PCTAIRE2
Epitope Specificity	301-400/523
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily.
SUBUNIT	Found in a complex containing CABLES1, CDK16 and TDRD7. Interacts with TDRD7
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with cyclins to phosphorylate key substrates involved in cell cycle progression. Another family of proteins, Cdk inhibitors, also play a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Members of the Cdk family include Cdk2-Cdk8, PCTAIRE-1-3, PITALRE and PITSLRE. PCTAIRE-1, PCTAIRE-2 and PCTAIRE-3 comprise a subfamily of cdc2-related serine/threonine kinases. PCTAIRE-1, which is expressed primarily in mammalian brain, interacts with a variety of proteins and is thought to be part of a multiple signal transduction cascade. PCTAIRE-2, also expressed in brain, may be important in terminally differentiated neurons.

Additional Information

Gene ID	5128
Other Names	PCTAIRE2; PCKT2; CDK17_HUMAN; CDK17; Cell division protein kinase 17; PCTAIRE-motif protein kinase 2; Serine/threonine-protein kinase PCTAIRE-2; 2.7.11.22; cyclin dependent kinase 17; PCTAIRE protein kinase 2
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CDK17
Synonyms	PCTAIRE2, PCK2
Function	May play a role in terminally differentiated neurons. Has a Ser/Thr-phosphorylating activity for histone H1 (By similarity).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.