

Neutrophil Elastase Rabbit pAb

Catalog # AP99102

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q3UP87
Other Accession	Q3UP87
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28648
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from mouse Neutrophil Elastase
Epitope Specificity	51-150/265
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the peptidase S1 family. Elastase subfamily. Contains 1 peptidase S1 domain.
SUBUNIT	Interacts with NOTCH2NL.
DISEASE	Defects in ELANE are a cause of cyclic haematopoiesis (CH) [MIM:162800]; also known as cyclic neutropenia. CH is an autosomal dominant disease in which blood-cell production from the bone marrow oscillates with 21-day periodicity. Circulating neutrophils vary between almost normal numbers and zero. During intervals of neutropenia, affected individuals are at risk for opportunistic infection. Monocytes, platelets, lymphocytes and reticulocytes also cycle with the same frequency. Defects in ELANE are the cause of neutropenia severe congenital autosomal dominant type 1 (SCN1) [MIM:202700]. SCN1 is a disorder of hematopoiesis characterized by a maturation arrest of granulopoiesis at the level of promyelocytes with peripheral blood absolute neutrophil counts below $0.5 \times 10^9/l$ and early onset of severe bacterial infections.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Elastases form a subfamily of serine proteases that hydrolyze many proteins in addition to elastin. Humans have six elastase genes which encode the structurally similar proteins. The product of this gene hydrolyzes proteins within specialized neutrophil lysosomes, called azurophil granules, as well as proteins of the extracellular matrix following the protein's release from activated neutrophils. The enzyme may play a role in degenerative and inflammatory diseases by its proteolysis of collagen-IV and elastin of the extracellular matrix. This protein degrades the outer membrane protein A (OmpA) of E. coli as well as the virulence factors of such bacteria as Shigella, Salmonella and Yersinia. Mutations in this gene are associated with cyclic neutropenia and severe congenital neutropenia (SCN). This gene is clustered with other serine protease gene family members, azurocidin 1 and proteinase 3 genes, at chromosome 19pter. All 3 genes are expressed coordinately and their protein products are packaged together into azurophil granules during neutrophil differentiation. [provided by RefSeq, May 2009].

Additional Information

Gene ID	50701
Other Names	ELA2; GE; HLE; HNE; NE; PMN-E; SCN1; F430011M15Rik; ELNE_HUMAN; ELANE; Bone marrow serine protease; Elastase-2; Human leukocyte elastase (HLE); Medullasin; PMN elastase; 3.4.21.37; ELNE_MOUSE; Leukocyte elastase; elastase, neutrophil expressed; elastase 2, neutrophil; neutrophil elastase; polymorphonuclear leukocyte elastase
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	Elane
Synonyms	Ela2
Function	Serine protease that modifies the functions of natural killer cells, monocytes and granulocytes (PubMed:10947984). Shows microbicidal activity in neutrophils by mediating cleavage and degradation of bacterial proteins (By similarity). Capable of killing E.coli but not S.aureus in vitro; digests outer membrane protein A (ompA) in E.coli and K.pneumoniae (PubMed:10947984). Involved for the formation of neutrophil extracellular traps (NETs): translocates into the nucleus downstream of MPO and mediates degradation of histone H4, thereby participating to chromatin decondensation (PubMed:20974816, PubMed:32045782). NETs are mainly composed of DNA fibers and are released by neutrophils to trap pathogens during inflammation (By similarity). Inhibits C5a-dependent neutrophil enzyme release and chemotaxis (By similarity). Promotes cleavage of GSDMB, thereby inhibiting pyroptosis (By similarity). Promotes blood coagulation (PubMed:20676107). Through the activation of the platelet fibrinogen receptor integrin alpha-IIb/beta-3, potentiates platelet aggregation induced by a threshold concentration of cathepsin G (CTSG) (By similarity). Cleaves and thus inactivates tissue factor pathway inhibitor (TFPI) (By similarity).
Cellular Location	Cytoplasmic vesicle, phagosome {ECO:0000250 UniProtKB:P08246}. Lysosome {ECO:0000250 UniProtKB:P08246}. Cytolytic granule {ECO:0000250 UniProtKB:P08246}. Nucleus {ECO:0000250 UniProtKB:P08246} Note=Localizes to azurophil granules, a type of primary lysosome found in neutrophils and monocytes that contains antimicrobial proteins (By similarity). Translocates from azurophil granules to the nucleus downstream of MPO during formation of neutrophil extracellular traps (NETs), and is present in released NETs (PubMed:32045782). Localized in phagolysosomes following ingestion of E.coli by neutrophils (By similarity). {ECO:0000250 UniProtKB:P08246, ECO:0000269 PubMed:32045782}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.