

LPA2 Rabbit pAb

Catalog # AP99242

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9HBW0
Other Accession	Q9HBW0
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38741
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human LPA2
Epitope Specificity	1-100/351
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the G-protein coupled receptor 1 family.
SUBUNIT	Interacts with SLC9A3R2/NHERF2, MAGI3 and PLCB3. Interacts with RALA and ADRBK1.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of family I of the G protein-coupled receptors, as well as the EDG family of proteins. This protein functions as a lysophosphatidic acid (LPA) receptor and contributes to Ca ²⁺ mobilization, a critical cellular response to LPA in cells, through association with Gi and Gq proteins. An alternative splice variant has been described but its full length sequence has not been determined. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	9170
Other Names	EDG-4; EDG4; LPA-2; LPA2; IPA2; LPAR2_HUMAN; LPAR2; LPA receptor 2; Lysophosphatidic acid receptor Edg-4; LPAR2_MOUSE; lysophosphatidic acid receptor 2; endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 4
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1 µg/Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	LPAR2 (HGNC:3168)
Synonyms	EDG4, LPA2
Function	Receptor for lysophosphatidic acid (LPA), a mediator of diverse cellular activities. Seems to be coupled to the G(i)/G(o), G(12)/G(13), and G(q) families of heteromeric G proteins. Plays a key role in phospholipase C-beta (PLC-beta) signaling pathway. Stimulates phospholipase C (PLC) activity in a manner that is independent of RALA activation.
Cellular Location	Cell surface. Cell membrane; Multi-pass membrane protein. Note=Prior to LPA treatment found predominantly at the cell surface but in the presence of LPA colocalizes with RALA in the endocytic vesicles
Tissue Location	Expressed most abundantly in testes and peripheral blood leukocytes with less expression in pancreas, spleen, thymus and prostate. Little or no expression in heart, brain, placenta, lung, liver, skeletal muscle, kidney, ovary, small intestine, or colon

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.