

CPEB4 Rabbit pAb

Catalog # AP99246

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q17RY0
Other Accession	Q17RY0
Host	Rabbit
Clonality	Polyclonal
Calculated MW	80152
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CPEB4
Epitope Specificity	301-400/729
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the RRM CPEB family. Contains 2 RRM (RNA recognition motif) domains.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	80315
Other Names	CPE-BP4; hCPEB-4; Cpeb-4; mCPEB-4; CPEB4_HUMAN; CPEB4; CPE-binding protein 4; KIAA1673; CPEB4_MOUSE; cytoplasmic polyadenylation element binding protein 4
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CPEB4
Synonyms	KIAA1673
Function	Sequence-specific RNA-binding protein that binds to the cytoplasmic polyadenylation element (CPE), an uridine-rich sequence element (consensus sequence 5'-UUUUUAU-3') within the mRNA 3'-UTR (PubMed: 24990967). RNA binding results in a clear conformational change analogous to the Venus fly

trap mechanism (PubMed:[24990967](#)). Regulates activation of unfolded protein response (UPR) in the process of adaptation to ER stress in liver, by maintaining translation of CPE- regulated mRNAs in conditions in which global protein synthesis is inhibited (By similarity). Required for cell cycle progression, specifically for cytokinesis and chromosomal segregation (PubMed:[26398195](#)). Plays a role as an oncogene promoting tumor growth and progression by positively regulating translation of t-plasminogen activator/PLAT (PubMed:[22138752](#)). Stimulates proliferation of melanocytes (PubMed:[27857118](#)). In contrast to CPEB1 and CPEB3, does not play role in synaptic plasticity, learning and memory (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q7TN98}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q7TN98}. Cell projection, dendritic spine {ECO:0000250|UniProtKB:Q7TN98}. Postsynaptic density {ECO:0000250|UniProtKB:Q7TN98}. Cell projection, axon {ECO:0000250|UniProtKB:Q7TN98}. Cell projection, growth cone {ECO:0000250|UniProtKB:Q7TN98}. Endoplasmic reticulum {ECO:0000250|UniProtKB:Q7TN98}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:Q7TN98}

Tissue Location

Expressed in pancreas in islets and ductal cells (at protein level) (PubMed:22138752). Expressed in melanocytes (PubMed:27857118).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.