10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

GAK Rabbit pAb

Catalog # AP99267

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	O14976
Other Accession	O14976
Reactivity	Rat, Mouse
Predicted	Human, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	143191
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human GAK
Epitope Specificity	331-430/1311
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the protein kinase superfamily. Ser/Thr protein kinase family.Contains 1 C2 tensin-type domain.Contains 1 J domain. Contains 1 phosphatase tensin-type domain. Contains 1 protein kinase domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Associates with cyclin G and CDK5. Seems to act as an auxilin homolog that is involved in the uncoating of clathrin-coated vesicles by Hsc70 in non-neuronal cells. Expression oscillates slightly during the cell cycle, peaking at G1. Tissue specificity: Ubiquitous. Highest in testis.

Additional Information

Gene ID	2580
Other Names	DNAJ26; DNAJC26; D130045N16Rik; GAK_HUMAN; GAK; DnaJ homolog subfamily C member 26; 2.7.11.1; GAK_MOUSE; cyclin G associated kinase; auxilin-2
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	GAK (HGNC:4113)

Function	Associates with cyclin G and CDK5. Seems to act as an auxilin homolog that is involved in the uncoating of clathrin-coated vesicles by Hsc70 in non-neuronal cells. Expression oscillates slightly during the cell cycle, peaking at G1 (PubMed:10625686). May play a role in clathrin-mediated endocytosis and intracellular trafficking, and in the dynamics of clathrin assembly/disassembly (PubMed:18489706).
Cellular Location	Cytoplasm, perinuclear region. Golgi apparatus, trans-Golgi network. Cell junction, focal adhesion. Cytoplasmic vesicle, clathrin-coated vesicle. Note=Localizes to the perinuclear area and to the trans-Golgi network. Also seen on the plasma membrane, probably at focal adhesions. Recruitment to clathrin- coated vesicles depends on temporal variations in phosphoinositide composition of clathrin-coated vesicles (PubMed:31962345)
Tissue Location	Ubiquitous. Highest in testis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.