

Ube2G1 Rabbit pAb

Catalog # AP99272

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P62253
Other Accession	P62253
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19509
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Ube2G1/UBC7
Epitope Specificity	101-170/170
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the ubiquitin-conjugating enzyme family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Ubiquitin is an abundant, highly conserved protein found in all eukaryotic cells either free or covalently attached to cellular proteins. The primary function of ubiquitin in mammalian systems is to clear abnormal, foreign, and improperly folded proteins by targeting them for proteasome degradation. In <i>Saccharomyces cerevisiae</i> , ubiquitin-like proteins include Rub1, Ula1, Uba3, Smt3, Ubc2, Ubc12 and Ubc9. Rub1 shares 53% homology with ubiquitin and requires activation via the E2 proteins, including Ula1, Uba3 and Ubc12 in order to conjugate to substrates directed to different proteolytic systems. Ubc4 catalyzes ubiquitination of IκBα in a phosphorylation and SCFB-TRCP dependent manner. In this particular reaction, E1 first transfers ubiquitin to the E2 component Ubc4, and Ubc4 then associates with E3 ligase, which conjugates the poly-ubiquitin chain on a target protein. In this fashion, the chain tags the IκBα for degradation by a proteasome thus lifting the inhibitory effect of IκBα on NFκB and allowing NFκB to enter the nucleus.

Additional Information

Gene ID	7326
Other Names	E217K; UBC7; UBE2G; 2700059C12Rik; D130023C12Rik; UB2G1_HUMAN; UBE2G1; E2 ubiquitin-conjugating enzyme G1; Ubiquitin carrier protein G1; Ubiquitin-protein ligase G1; 2.3.2.23; UB2G1_MOUSE; ubiquitin conjugating enzyme E2 G1; ubiquitin-conjugating enzyme E2G 1 (homologous to <i>C. elegans</i> UBC7); ubiquitin-conjugating enzyme E2G 1 (UBC7 homolog, <i>C. elegans</i>); ubiquitin-conjugating enzyme E2G 1 (UBC7 homolog, yeast); ubiquitin-conjugating enzyme E2G 1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	UBE2G1
Synonyms	UBE2G
Function	Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'Lys-48', as well as 'Lys-63'-linked polyubiquitination. May be involved in degradation of muscle-specific proteins. Mediates polyubiquitination of CYP3A4.
Tissue Location	Widely expressed, mainly in skeletal muscle.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.