

AMH Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9940C

Product Information

Application	WB, FC, IHC-P, E
Primary Accession	P03971
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	59195
Antigen Region	424-451

Additional Information

Gene ID	268
Other Names	Muellerian-inhibiting factor, Anti-Muellerian hormone, AMH, Muellerian-inhibiting substance, MIS, AMH, MIF
Target/Specificity	This AMH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 424-451 amino acids of human AMH.
Dilution	WB~~1:1000 FC~~1:25 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	AMH Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	AMH (HGNC:464)
Synonyms	MIF
Function	Plays an important role in several reproductive functions. Induces Muellerian duct regression during male fetal sexual differentiation (PubMed: 34155118 , PubMed: 3754790 , PubMed: 8469238). Also plays a role in


Leydig cell differentiation and function (By similarity). In female acts as a negative regulator of the primordial to primary follicle transition and decreases FSH sensitivity of growing follicles (PubMed:[14742691](#)). AMH signals by binding to a specific type- II receptor, AMHR2, that heterodimerizes with type-I receptors (ACVR1 and BMPR1A), and recruiting SMAD proteins that are translocated to the nucleus to regulate target gene expression (PubMed:[20861221](#), PubMed:[34155118](#)).

Cellular Location	Secreted
Tissue Location	In ovaries, AMH is detected in granulosa cells of early growing follicles.


Background

Anti mullerian hormone (AMH) is a member of the TGF beta superfamily. It is secreted as a homodimeric 140kD disulphide linked precursor that is cleaved to release the mature 30kD homodimer. Originally classified as a foetal testicular hormone that inhibits Mullerian duct development, AMH is expressed post natally by immature Sertoli cells, and to a lesser degree by granulosa cells. AMH plays a role in testicular differentiation and in the regulation of ovarian follicle growth.

Images




All lanes : Anti-AMH Antibody (Center) at 1:1000-1:2000 dilution Lane 1: Human cerebellum lysate Lane 2: HeLa whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: SH-SY5Y whole cell lysate Lane 5: U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 59 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AMH Antibody (Center) (Cat. #AP9940c) IHC analysis in formalin fixed and paraffin embedded prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the AMH Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

AMH Antibody (Center) (Cat. #AP9940c) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Citations

- [FOXL2 Is an Essential Activator of SF-1-Induced Transcriptional Regulation of Anti-Müllerian Hormone in Human Granulosa Cells.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.