

Tal1 Rabbit pAb

Catalog # AP99542

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	P17542
Other Accession	P17542
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34271
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Tal1
Epitope Specificity	185-300/331
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 basic helix-loop-helix (bHLH) domain.
SUBUNIT	Efficient DNA binding requires dimerization with another bHLH protein. Forms heterodimers with TCF3. Binds to the LIM domain containing protein LMO2 and to DRG1. Can assemble in a complex with LDB1 and LMO2. Component of a TAL-1 complex composed at least of CBFA2T3, LDB1, TAL1 and TCF3.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Activation of TAL1 characterizes up to 60% of cases of human T cell acute lymphoblastic leukemia, making it the most frequent gain-of-function mutation observed in this disorder. TAL1 (also designated SCL) is a serine phosphoprotein and basic helix-loop-helix transcription factor known to regulate embryonic hematopoiesis. This transcription factor binds as a heterodimer with E2A and HEB/HTF4 to a nucleotide sequence motif termed the E-box. In addition, leukemogenesis is accelerated dramatically by transgenic co-expression of TAL1 and the catalytic subunit of casein kinase IIalpha, a serine/ threonine protein kinase known to modulate the activity of other bHLH transcription factors.

Additional Information

Gene ID	6886
Other Names	SCL; TCL5; bHLHa17; tal-1; TAL1_HUMAN; TAL1; Class A basic helix-loop-helix protein 17 (bHLHa17); Stem cell protein; T-cell leukemia/lymphoma protein 5; TAL bHLH transcription factor 1, erythroid differentiation factor; T-cell acute lymphocytic leukemia 1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TAL1
Synonyms	BHLHA17, SCL, TCL5
Function	Implicated in the genesis of hemopoietic malignancies. It may play an important role in hemopoietic differentiation. Serves as a positive regulator of erythroid differentiation (By similarity).
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00981}.
Tissue Location	Leukemic stem cell.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.