

IQGAP1 Recombinant Rabbit mAb

Catalog # AP99553

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P46940
Other Accession	P46940
Host	Rabbit
Clonality	Recombinant
Calculated MW	189252
Physical State	Liquid
Immunogen	A synthesized peptide derived from human IQGAP1
Epitope Specificity	1-58/1657
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol.
SIMILARITY	Contains 1 CH (calponin-homology) domain. Contains 4 IQ domains. Contains 1 Ras-GAP domain. Contains 1 WW domain.
SUBUNIT	Interacts with CDC42; the interaction is demonstrated with IQGAP1 in GTP-bound and in nucleotide-free state. Interacts with RAC1. Does not interact with RHOA. Interacts with TSG101. Interacts with PAK6.
Post-translational modifications	Phosphorylation of Ser-1443 by PKC/PRKCE prevents interaction between C1 and C2, allowing binding of nucleotide-free CDC42. Ser-1443 phosphorylation enhances the ability to promote neurite outgrowth.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Binds to activated CDC42 but does not stimulate its GTPase activity. It associates with calmodulin. Could serve as an assembly scaffold for the organization of a multimolecular complex that would interface incoming signals to the reorganization of the actin cytoskeleton at the plasma membrane.

Additional Information

Gene ID	8826
Other Names	HUMORFA01; SAR1; p195; D7Ert237e; D7Ert257e; mKIAA0051; IQGA1_HUMAN; IQGAP1; KIAA0051; IQGA1_MOUSE; IQ motif containing GTPase activating protein 1; RasGAP-like with IQ motifs
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:50-200,IF=1:100-500,Flow-Cyt=1:50-100
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	IQGAP1
Synonyms	KIAA0051
Function	Plays a crucial role in regulating the dynamics and assembly of the actin cytoskeleton. Recruited to the cell cortex by interaction with ILK which allows it to cooperate with its effector DIAPH1 to locally stabilize microtubules and allow stable insertion of caveolae into the plasma membrane (By similarity). Binds to activated CDC42 but does not stimulate its GTPase activity. Associates with calmodulin. May promote neurite outgrowth (PubMed:15695813). May play a possible role in cell cycle regulation by contributing to cell cycle progression after DNA replication arrest (PubMed:20883816).
Cellular Location	Cell membrane. Nucleus. Cytoplasm. Cytoplasm, cell cortex {ECO:0000250 UniProtKB:Q9JKF1}. Apical cell membrane. Basolateral cell membrane {ECO:0000250 UniProtKB:Q9JKF1}. Note=Subcellular distribution is regulated by the cell cycle, nuclear levels increase at G1/S phase (PubMed:20883816).
Tissue Location	Expressed in the placenta, lung, and kidney (PubMed:8051149). A lower level expression is seen in the heart, liver, skeletal muscle and pancreas (PubMed:8051149)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.